

## PERFORMANCE OUTCOMES BENCHMARKING

### CROSS-REFERENCE TO RELATED APPLICATIONS

This patent application claims priority to U.S.  
Provisional Patent Application No. 60/252,129, filed

5 November 21, 2000, which is hereby incorporated by reference  
herein in its entirety.

### FIELD OF THE INVENTION

10 The present invention relates generally to creation and  
management of feedback reports for a benchmarking process  
and, more particularly, to a technique for acquiring primary  
data at the source, compiling the data in an analytically  
meaningful manner for benchmarking, preparing web based  
reports and maintaining and managing cumulative historical  
15 data and reports.

### BACKGROUND OF THE INVENTION

Benchmarking which is a comparison of performance  
outcomes from a process to a norm is a useful tool for  
20 measuring quality and developing methods of improving a  
process.

One example of an area in which benchmarking is  
particularly useful is in monitoring ambulatory surgery

procedures. Professional organizations have made attempts  
to acquire and tabulate ambulatory surgical procedures  
outcomes data for benchmarking. These attempts have  
suffered three deficiencies. First the data is typically  
5 taken from secondary sources such as billings or claims  
records and not from the patient. Second, uniform criteria  
are not applied in obtaining the data collection. Third,  
data is typically compiled and distributed in a hard copy  
form and cannot be accessed in a convenient internet  
10 dispersed form.

#### SUMMARY OF THE INVENTION

In view of the foregoing it would be desirable to  
provide a technique for collecting, evaluating and reporting  
outcomes data for the purposes of benchmarking which  
15 overcomes the above described inadequacies and shortcomings.  
More particularly it would be desirable to provide a  
technique for collecting outcomes data at a primary level,  
evaluating the data in an analytically meaningful manner  
reporting the outcome data and maintaining historical data  
20 for generation of future comparisons in an efficient and  
cost effective manner. It is particularly desirable that  
such a system be computer based and use a network such as  
the internet both for transporting input data and  
disseminating reports.

According to the present invention, a technique for performance outcomes benchmarking is provided. In one exemplary embodiment, the technique is realized by a method for outcomes monitoring comprising the steps of: collecting  
5 at least two outcomes data sets; converting the at least two outcomes data sets into an at least one outcomes result; establishing a norm for an outcomes data group, the outcomes data group comprising a plurality of the at least two outcomes data sets; comparing a selected one of the at  
10 least one outcomes result to the norm; and generating at least one outcomes monitoring report comprising the selected one of the at least one outcomes result and the norm.

In accordance with other aspects of this exemplary embodiment, the method may further include: transmitting  
15 the at least two outcomes data sets to a data processor; selectively restricting access to the outcomes monitoring report; posting the outcomes monitoring report to a webpage or selectively restricting access to the webpage or a combination thereof.

20 In accordance with other aspects of this exemplary embodiment, the outcomes data sets may be collected from at least one user entity at a plurality of discrete intervals and the outcomes report may be prepared from the outcomes

data collected at least two of the plurality of discrete intervals.

In accordance with further aspects of this exemplary embodiment, the outcomes data sets may be collected from a plurality of user entities; outcomes data sets for each user entity of the plurality of user entities individually identified and converted. The outcomes data sets from the plurality of user entities may comprise the outcomes data group.

In accordance with further aspects of this exemplary embodiment the outcomes monitoring report includes at least one outcomes result for a selected user entity of the plurality of user entities and at least one comparison of the norm to the selected one of the at least one outcomes result for the selected user entity.

In accordance with additional aspects of this exemplary embodiment, the embodiment includes a computer signal embodied in a carrier wave readable by a computing system and encoding a computer program of instructions for executing a computer process performing the method for outcomes monitoring described herein.

In another exemplary embodiment a technique for monitoring surgical procedure outcomes is provided. In one exemplary embodiment the technique is realized by a method

for outcomes monitoring of surgical procedures comprising  
the steps of: collecting at least two primary source  
surgical outcomes data sets; converting the at least two  
primary source surgical outcomes data sets into at least one  
5 outcomes result; establishing a norm for an outcomes data  
group, the outcomes data group comprising a plurality of the  
at least two outcomes data sets, comparing a selected one of  
the at least one outcomes result to the norm; and generating  
at least one outcomes monitoring report comprising the  
10 selected one of the at least one outcomes result and the  
norm.

In accordance with other aspects of this exemplary  
embodiment the method may further include: transmitting the  
at least two primary source surgical outcomes data sets to a  
15 data processor; selectively restricting access to the  
outcomes monitoring report; posting the outcomes monitoring  
report to a webpage; and selectively restricting access to  
the webpage.

In accordance with other aspects of this exemplary  
20 embodiment the method may further include collecting the at  
least two primary source surgical outcomes data sets from a  
plurality of surgical centers; and individually identifying  
and converting the at least two primary source outcomes data  
sets for each surgical center of the plurality of surgical

099645-12001  
FOOT 58367.000003

centers where the outcomes data sets from the plurality of surgical centers comprises the outcomes data group.

In accordance with other aspects of this exemplary embodiment the method may further include an outcomes  
5 monitoring report which has at least one outcomes result for a selected surgical center of the plurality of surgical centers and at least one comparison of the norm to the selected one of the at least one outcomes result for the selected surgical center.

10 In accordance with further aspects of this exemplary embodiment, the embodiment may include a computer signal embodied in a carrier wave readable by a computing system and encoding a computer program of instructions for executing a computer process performing the method for  
15 outcomes monitoring of surgical procedures described herein.

In another exemplary embodiment an apparatus for outcomes monitoring is provided. The apparatus comprises: a data collection portion wherein the data collection portion collects at least two outcomes data sets; a data processor  
20 portion wherein the data processor portion receives the at least two outcomes data sets from the data collection portion and wherein the data processor comprises: a converter portion wherein the converter portion converts the at least two outcomes data sets into an at least one

099647-13001  
58367.000003

outcomes result; a norm establishing portion wherein the  
norm establishing portion establishes a norm for an outcomes  
data group, the outcomes data group comprising a plurality  
of the at least two outcomes data sets, a comparison portion  
5 wherein the comparison portion compares a selected one of  
the at least one outcomes result to the norm; and a report  
generation portion wherein the report generation portion  
generates at least one outcomes monitoring report comprising  
the selected one of the at least one outcomes result and the  
10 norm.

In accordance with further aspects of this exemplary  
embodiment, the apparatus for outcomes monitoring may  
further comprise a webpage portion wherein the at least one  
outcomes monetary report is posted to a webpage; or a  
15 security portion, the security portion selectively  
restricting access to the at least two outcomes data sets,  
the at least one outcomes result and the at least one  
outcomes monitoring report; or both.

In accordance with other aspects of this exemplary  
20 embodiment, the apparatus for outcomes monitoring may be an  
apparatus wherein the at least two outcomes data sets are  
surgical procedures outcomes data sets, or wherein the at  
least two surgical procedures outcomes data sets are primary  
source data sets or both.

09996475-112001

The present invention will now be described in more detail with reference to exemplary embodiments thereof as shown in the appended drawings. While the present invention is described below with reference to preferred embodiments, 5 it should be understood that the present invention is not limited thereto. Those of ordinary skill in the art having access to the teachings herein will recognize additional implementations, modifications, and embodiments, as well as other fields of use, which are within the scope of the 10 present invention as disclosed and claimed herein, and with respect to which the present invention could be of significant utility.

#### BRIEF DESCRIPTION OF THE DRAWINGS

15 In order to facilitate a fuller understanding of the present invention, reference is now made to the appended drawings. These drawings should not be construed as limiting the present invention, but are intended to be exemplary only.

20 Figure 1 is a schematic diagram of the system of the invention in accordance with the present invention;

Figure 2 is a diagram showing the steps of using the invention in accordance with one embodiment of the present invention;



Figure 3 is a exemplary up load screen in accordance with one embodiment of the invention;

Figure 4 is an exemplary report in accordance with one embodiment of the invention;

5        Figure 5 shows a flow chart for the report system structure in accordance with one embodiment of the invention;

10        Figure 6 shows exemplary parameters for a file for defining report periods in accordance with one embodiment of the invention;

Figure 7 shows exemplary parameters for a file which provides paths to different components of a report system in accordance with one embodiment of the invention;

15        Figure 8 shows an example of the parameters associated with an executive table list file in accordance with one embodiment of the invention;

Figure 9 show an example of a data table items file parameters in accordance with one embodiment of the invention;

20        Figure 10 shows an example of a data table in accordance with one embodiment of the invention;

Figure 11 shows an example of parameters for general report sections in accordance with an embodiment of the invention;

Figure 12 shows an example of parameters of an indicators description string in accordance with an embodiment of the invention;

Figure 13 shows an example of an indicator description report page in accordance with one embodiment of the invention;

Figure 14 shows an example of parameters of a comparison table file in accordance with one embodiment of the invention;

Figure 15 shows a report of the type showing data for a user and cumulative data for all user groups used in establishing a norm in accordance with one embodiment of the invention;

Figure 16 shows the parameters of a ProcDistrib file in accordance with one embodiment of the invention;

Figure 17 shows a case distribution table in accordance with one embodiment of the invention;

Figure 18 shows an exemplary first stage list file in accordance with one embodiment of the invention;

Figure 19 shows an exemplary second stage list file in accordance with one embodiment of the invention;

Figure 20 shows an exemplary report folder structure for a private web site in accordance with one embodiment of the invention;

Figure 21 is an exemplary menu for accessing previous reports in accordance with one embodiment of the invention;

Figure 22 is an exemplary graphic report in accordance with one embodiment of the invention; and

5        Figure 23 in an exemplary graphic report in accordance with one embodiment of the invention.

#### DETAILED DESCRIPTION OF EXEMPLARY EMBODIMENTS

0996475-12004  
10021 5249550

The outcomes monitoring system (OMS) of the invention  
10    (also referred to herein as the System) is described in the context of collecting and reporting outcomes data for benchmarking ambulatory surgical procedures which may be used for benchmarking both a given center's performance for a particular procedure(s) against historical data for that  
15    surgery center or for benchmarking the performance of a particular center with respect to the performance of a group of centers. As used herein a center is a user unit entity such as a clinic, or a business for example. The term "user" includes both the user entities as well as the human  
20    beings using the system. As one skilled in the art will appreciate this is one example of the usefulness of the invention. The invention may be applied in other areas of healthcare, such as, the benchmarking of hospital, nursing

Further the invention may be generally applicable to benchmarking performance in such areas as sales force

performance, customer satisfaction, manufacturing process performance, service provider performance and the like. As one of ordinary skill in the art will appreciate the outcome monitoring system of the invention may be used for benchmarking performance in these areas and other areas without undue experimentations. Further, the invention accommodates benchmarking from primary data (e.g., information directly acquired from individuals involved in the process such as patients, health care providers, or customers and the like or measurements made or data taken as a direct result of the process). For example, or secondary data such as billing records or claims records, for example, or a combination thereof. The ability to obtain benchmarking information from a primary source while maintaining confidentiality or privacy of specific identifiers for the primary source is a particularly desirable feature of some embodiments of the invention.

At this point it should be noted that the system in accordance with the present invention as described herein typically involves the processing of input data and the

generation of output data to some extent. This input data processing and output data generation may be implemented in hardware or software. For example, specific electronic components may be employed in a personal computer, server or similar or related circuitry for implementing the functions associated with acquiring, transmitting, analyzing and storing data in accordance with the present invention as described herein. Alternatively, one or more processors operating in accordance with stored instructions may implement the functions associated the system in accordance with the present invention as described herein. If such is the case, it is within the scope of the present invention that such instructions may be stored on one or more processor readable media, or transmitted to one or more processors via one or more signals.

Referring to Figure 1, in general the system of the invention comprises a plurality of user interfaces 500, 520, 540, 560, a network 100 and a server 200. Four user interfaces are shown in Figure 1 as representative of the plurality of user interfaces that may be associated with the system. The number of user interfaces may be greater or less than four. The plurality of user interfaces are personal computers (PC) equipped with modems and suitable software in an exemplary embodiment. More particularly

5 suitable user interfaces 500, 520, 540, 560 may, for  
example, include an IBM-PC compatible personal computer  
capable to run under Windows-95 (or 98) operating system, a  
Pentium processor with clock speed more than 100 MHz, 16  
10 MBytes (or higher) of RAM, 1000 MByte (or higher) hard disk  
capacity; a dial-up Internet connection with an Internet  
service provider; a 28.8 (or higher) kbps telephone modem  
for data transfer; (optionally) a lower speed modem for  
automated patient interview dialing, which requires a second  
10 telephone line; and the database management system such as  
FileMaker Pro or Visual Basic.

15 Additionally, the user interfaces 500, 520, 540, 560  
will have performance outcomes benchmarking software as  
described herein. Further, it is desirable that the user  
15 interfaces 500, 520, 540, 560 have a means for archiving  
data such as a floppy disk or CD or the like or be linked to  
a suitable memory device for archiving data obtained at the  
user site.

20 The user interfaces 500, 520, 540, 560 are used for  
collecting data and inputting data into the system and  
accessing reports generated by the system. Further, as  
illustrated in Figure 1 the user interfaces 500, 520, 540,  
560 are not directly interconnected. This is preferable in  
embodiments where highly confidential information is

involved such as when the users are competitors or when  
personal confidential information such as patient identity  
is involved, for example. In such cases, it is important to  
restrict access of a particular user to that user's data and  
5 results, and that user's results as compared to compiled  
benchmarking norms for all users.

As Figure 1 shows the user interfaces 500, 520, 540,  
560 are connected to a network 100. The network 100 is used  
to transmit data collected and input by the users to the  
10 server and disseminate the analyzed data and reports to the  
user. In an exemplary embodiment, the network 100 is a  
general access network such as the Internet, for example.  
Confidentiality of information transmitted over the network  
100 may be maintained by the use of such known means as  
15 password protection and encryption, for example. In  
preferred embodiments reports are posted to a password  
protected web page or the like.

The server 200 processes the collected data transmitted  
via the network 100, prepares reports and archives data and  
20 reports. The algorithms of this invention typically reside  
in the server 200. The algorithms are applied to analyze  
the collected data which in turn yields results suitable for  
comparisons of performance. The comparisons may be internal  
to a particular user such as comparison of the performance

of one month to the performance of another month, for example, or may compare the user's performance for a specified period to a norm based on an average for a group of users. The results may be considered to be statistically significant if based on a data group (as used herein a data group is a plurality of discrete data sets) of a suitable size. Typically, at least 30 data points are used for calculating statistically significant results. Reports reflecting the results of the data analysis are generated in the server 200. In an exemplary embodiment results are stored in tables such that reports customized in content and form may be generated in a facile manner. The collected data, analyzed results and reports or a combination thereof are also archived in the server 200 or alternatively in a memory device associated with the server. Hence not only may previous reports be accessed, but also collected data and analyzed results are available for use in generating future reports. For example it is frequently desirable to compare existing data with new data as it becomes available.

In an exemplary embodiment, the server used to operate an Outcomes Monitoring System may be a Hewlett Packard NetServer LH 3. The HP Server contains 5 hard drives which are used for the storage of data and the programs required for operation. In conjunction with the HP Server, a Cisco



1720 Router is used to provide the Internet connection.

Internet service with a connection speed of 512K is used in this exemplary embodiment. This server is described for illustrative purposes and is one of many servers suitable

5 for use in the outcomes monitoring system of this invention.

Referring to Figure 2, use of the invention in an embodiment directed to outcomes monitoring of ambulatory surgical procedures, is summarized. As one skilled in the art will recognize this is one of many possible embodiments  
10 related to health care outcomes monitoring. Further, other embodiments of the invention may be applied in many other types of outcomes monitoring of processes including, for example, service performance, sales force performance, service provider performance, manufacturing performance and  
15 the like, for example. For convenience, use of the invention is described in terms of outcomes monitoring of ambulatory surgical procedures with the understanding that the fundamental steps of use of the invention are comparable in other embodiments.

20 As shown in Figure 2, the steps of using an exemplary embodiment of the invention directed to outcomes monitoring of ambulatory surgical procedures include: collection of patient, procedure and outcomes data S200, input of collected data S300, transmission of data to the system

server S400, compilation and analysis of data by system  
server algorithms S500, preparation of outcomes monitoring  
(benchmarking) reports by the system server S600, posting of  
the outcomes monitoring reports to a web page S700, and user  
5 review of reports via internet connection to the web page  
S800. The steps for use of the exemplary embodiment are  
discussed in further detail below.

Referring to Figure 2, step S200 comprises collection  
of patient, procedure and outcomes data (herein collectively  
10 referred to as outcomes data set(s)). In a preferred  
embodiment, the data is collected directly from the patient,  
center staff or both at a time coinciding with the  
procedure, within a predetermined proximity to the procedure  
or both.

15 To maintain patient privacy and confidentiality,  
patient identification codes may be assigned. A system of  
patient identification (ID) codes, unique for each episode  
of care, may be established at each center. In order to  
maintain complete patient confidentiality, it is preferable  
20 that centers not use social security numbers for ID codes.  
A surgical center may assign a patient a facility specific  
ID number, with no relation to any personal identification  
numbers or names. Alternatively, the ID "number" may be  
derived from another numbering system used by the facility,

such as patient account numbers, etc. so long as numbers are not repeated for patients with multiple episodes of care.

To protect privacy and confidentiality it is preferable that patient names and medical record numbers are not provided to

5 the server. Maintaining records of specific patient information associated with data entries at a center or local user level is important to the center's ability to address issues identified in the reports of the OMS and apply benchmarking results to improve performance.

10 Potential OMS participants in the exemplary embodiment are all patients scheduled for an applicable procedure. Participants as used herein are primary data sources. Criteria may be established to identify a suitable primary data source. For example patients in an exemplary  
15 embodiment for ambulatory surgical procedure monitoring are typically excluded if

- they undergo two unrelated procedures simultaneously (example: patient having a cataract removal and hernia repair at the same  
20 time);
- the patient is scheduled for a planned inpatient admission postoperatively; or

- none of the procedure codes on the patient's record matches any of the codes included in the identified procedure codes to be benchmarked.

5 In an exemplary embodiment, centers are strongly encouraged to submit data on as many patients as possible, (20 percent of specified population(s) is suggested) but there are no minimum requirements for data submission. Depending upon the size of the center and the volume of  
10 procedures performed, a center may determine its own standards for quantity of data collected. It should be kept in mind, however, that larger databases provide more meaningful information. A minimum of 30 points are desirable for statistically significant evaluation.  
15 However, lesser numbers may be useful for identifying quality issues.

Further the system permits benchmarking of a wide range of types of data. Hence many kinds a data may be collected and considerably flexibility is afforded in outcomes that  
20 may be monitored and benchmarked. However, for benchmarking a particular outcome it is important to use uniform criteria when collecting data related to that outcome.

In an embodiment for monitoring health care outcome stripping all personal identifiers from the data transmitted

to the server is desirable. It may further be desirable to inform patients that:

1. no identifiable medical or personal information will leave the Local Center;
2. that server staff will have no personal identifiers;
3. that only grouped information will be reported back to the center;
4. that participation will not affect the care they receive; and the like.

Additionally, in some cases it may be desirable to obtain informed consent from patient.

Referring again to Figure 2, step S300 is data input. The data collected will typically be collected as a data set for a particular process such as a surgical procedure, for example. Typically, a data set will include a plurality of responses to a set of indicators. Indicators may be verbal responses; measured analytical data such as times, weights, ages, dosage of medicine and the like, or observations of a third-party observer, for example. Any indicator reflective of the outcome of the process being monitored may be used. Although one indicator may be used, typically a set of indicators will be used. It may be desirable in some embodiments to validate benchmarking indicators utilizing

statistical methods known to those of ordinary skill in the art.

The indicators (indicators as used herein are the specific information gathered) may be defined for any process or procedure to be benchmarked. Such indicators may be customized to the procedure or process to be benchmarked.

The specific data collected for comparison purpose should be collected in a standardized manner with specified collection criteria. However, the system provides extensive flexibility in selection of data to be collected. This allows the system to be applied to a range of needs and benching marking of many types of procedures. Further data collection may include questions which are intended for site specific benchmarking in addition to questions which are intended for broader based comparisons. Additionally, questions for general reference such as marketing studies, scientific research studies and the like, may be included and archived for reference, but not utilized in the process of evaluating data and preparing benchmarking reports.

The data collection process may take many forms. The following example is illustrative of the type of information that may be collected and a procedure that may be followed which utilizes standardized criteria. As one skilled in the art will recognize, this is one of many suitable data

collection protocols which may be used in the practice of  
the invention.

Data collection in an Outcomes Monitoring System for  
Ambulatory Surgical Centers in one embodiment may begin by

5 opening a Medical Record Abstract Form. The form may  
include input fields such as:

- Add A New Record
- Patient ID
- Procedure Date
- 10 • Payor
- Procedure Information(predefined procedure  
category, codes facilitate standardization)
- Start and end time of the Procedures
- Recovery Sites
- 15 • Discharge Time
- Time at which Patient Met Discharge Criteria
- Type of Anesthesia (Codes facilitate  
standardization)
- MD Present during anesthesia (yes or no)
- 20 • Patient Disposition following surgery
- Any of a predetermined list of identified problems  
experienced (problems are defined as well as

criteria for assessing significance of a problem  
and response selections provided)

- Postoperative Pain Management (Including yes/no  
responses to questions such as:

5                   Pain Verbalized?

                  Med Ordered?

                  Med Admin?

                  and Pain Relieved?

- Discharge Pain Management (including yes/no  
10                   questions such as):

                  Pain Prescription Given?

                  Pain Control Methods Explained?:

                  and Institution Specific Variables

As a general rule, it is desirable for monitoring  
15   outcomes of ambulatory surgery procedures that a post-  
discharge patient interview take place within 1 - 2 working  
days from the date of discharge. In an exemplary embodiment  
this may be a phone interview. It is important that  
patients be contacted while the events of their care are  
20   fresh in their memory. The purpose of the interview is to  
collect information about the "outcome" of the procedure  
(i.e. checking the patient's condition after discharge), and  
to collect information about the patient's perception of



5

10

15

20

2. At any time after leaving the center did you have...

These questions relate to Quality - "Patients

experiencing problems after discharge related to the surgical procedure requiring medical or surgical care." The "key" words in this indicator are related to the surgical procedure and requiring medical or surgical care. Problems patients report that do not relate to their surgical procedure (i.e. a fall after arriving home; an auto accident on the way home, etc.) should not be considered in question 2 of the interview. Also, only problems that resulted in the patient seeking additional medical care may be considered as "meeting" the indicator. Questions applicable to specific procedure groups may be included.

3. After returning home... These questions relate to whether - Patients expressing pain after discharge who had relief of pain after utilizing pain control methods as instructed.

4. How well would you say your pain was relieved...

This question assess the patients' relief of pain. Patients may be asked to rate their pain on a scale of 1 to 10, with "1" equaling complete relief and "10" not relieved at all.

5. How would you describe the quality of care you received. . . . These questions are directed towards determining whether - patients were satisfied with pre-operative, intraoperative, and postoperative care.

For an exemplary embodiment for monitoring ambulatory surgical procedures, outcomes procedures to be monitored and indicators are preferably determined prior to data collection.

In an embodiment for benchmarking ambulatory surgical procedures, patient data is typically collected from patients, patient relatives or surgery center staff or a combination thereof.

In a preferred embodiment electronic forms and tables may be used to enter and compile patient data. As discussed each patient receives a unique ID code, so that data entered at a center may be transmitted to the server without personal identifiers.

Data may be entered directly into electronic screens during medical record reviews and telephone interviews. Alternatively data may be documented as paper records and entered into the electronic system at a later time. Data entry directly from a primary data source into electronic form is preferred. Further in a preferred embodiment data

collection forms are based on modifications of instruments that have been validated.

More particularly, data collection and abstracting may be completed in one of two ways:

- 5           1.   Manually: Blank forms are printed and data is  
                  manually entered onto the forms and entered into  
                  the computer file at a later time. This method is  
                  not generally recommended but can be used.
2.   Directly: Data are entered directly into the  
10               computer screen. After a sufficient number of  
                  records are completed (20 or more, for example),  
                  the data may be "uploaded" to server through an  
                  Internet connection.

15           As a center compiles and transmits information, server  
             staff may conduct both overall and center-specific analyses  
             of the clinical data. A center is encouraged to archive all  
             patient data with patient identifiers locally (See S320).  
             Such local archival enables a center to investigate issues  
20           identified by the OMS in depth at a later date, e.g.,  
             benchmarking identifies trends and follow-up of individual  
             patients may be necessary to establish reasons for  
             deviations from expected values. Thus by monitoring patient  
             specific data at the local level this information which is

useful to the center is maintained while the risk of  
improper invasion of patient privacy is minimized.

Because medical information is rightly held by patients  
and health care providers to be confidential, effort should  
5 be made to assure confidentiality and privacy throughout the  
system. Entry into the database associated with data input  
may be controlled via passwords for example, and only  
personnel directly involved in data entry or supervision  
should have access to databases to maintain confidentiality  
10 of patient data.

Other measures to restrict access to the information  
contained in the OMS may include physical and electronic  
barriers to access, restrictions on patient information  
maintained at the server and reporting only of group  
15 results. In embodiments in which data may be stored on the  
server, the server may be password protected at the user  
level, and a second password may be required to access the  
study database. Further in a preferred embodiment the  
personal identifiers are stripped from the data at the local  
20 level so that the system data base contains no personal  
identifiers.

The data may be collected by a designated individual or  
collected by various individuals during the course of the  
procedure and follow-up data collection. In some exemplary

embodiments collection by various individuals may be desirable to facilitate acquiring data in the optimum time frame with respect to performance of the procedure.

As shown in Figure 2, in step S400, the input data is transmitted to the server. In some exemplary embodiments, it is desirable to strip the data of any confidential identifiers prior to transmission. Further in some embodiments it is preferable to batch data prior to transmission to the server and use a File Transfer Protocol (FTP) to send data to the server.

In an exemplary embodiment the transmission process may begin by establishing a network connection to the server and selecting a software option for transmission, such as an "upload" icon, for example. An upload screen will then appear, with a series of steps. As each step is successfully completed, the next step will appear and the process may be completed by simply following each step in the program. An exemplary upload screen is shown in Figure 3.

In the exemplary embodiment shown in Figure 3, the second step verifies that the user's Internet connection is stable and will accommodate a smooth transfer of data. The user may also be asked for a password during this process.

The third step initiates the actual transfer (or upload) of data from the center to the system server. The user may be asked for a password during this process.

The fourth step in completing data transfer in the exemplary embodiment of Figure 3 is to move all files into the "stock" file. The stock file is the name of the folder where each individual record is stored. This permits later access to individual records or data taken for information only. Such information is typically stored locally (e.g., the user site) to maintain confidentiality yet provide necessary information for follow-up of issues identified in benchmarking. The "stock file" transfer procedure in one embodiment is the archiving of data locally as shown in step S320 of Figure 2.

Again, referring to Figure 3, the upload program may be exited upon confirmation of a successful transfer of data as indicated by the software.

Referring again to Figure 2 in step S500, the system server algorithms compile and analyze input data. Exemplary performance monitoring software is described in detail below. In general, the software has the capacity to convert an outcomes data set to an outcomes result. For example, data for a particular procedure may be selected from data for all procedures; data for a particular indicator for a

particular procedure accumulated during a specific time  
period may be averaged, and the like. Outcomes results may  
be derived from direct manipulation of input data (e.g.,  
outcomes data set), manipulation of other outcomes results,  
5 or any combination thereof.

The performance monitoring software can use multiple  
outcomes data sets to establish a norm for an outcomes data  
group. For example, in the case of ambulatory surgery  
centers, the data from all centers or a selected group of  
10 centers may be averaged to determine a norm. Alternatively,  
a norm for a single center may be established by averaging  
outcomes data sets for that center taken at designated time  
intervals, for example.

Benchmarking is achieved by comparing an outcomes  
15 result for an indicator or procedure or the like to the norm  
for that indicator or procedure. As will be apparent in the  
detailed discussion of the exemplary embodiment of the  
software of the invention, considerable customization of  
specific benchmarking is permitted within the system of the  
20 invention.

Referring again to Figure 2, step S600 is the  
production of the product of the benchmarking process or the  
feedback report(s) to the user (also referred to herein as  
outcomes monitoring reports, outcomes reports or reports).



0996475-12001  
Docket 52496660

In an Exemplary embodiment reports include benchmarking in which data collected through the system is analyzed to reflect compliance with each of a predetermine number of quality indicators. Although each center has access only to their own data and compliance rates, the reports also may show "average" compliance rates across all participating centers. Individual centers can then compare their rate of compliance against an overall average for a group of centers for each quality indicator or compare an individual center's compliance to that individual center's compliance for another period. Password protection may be used to appropriately restrict access to reports. In one preferred embodiment, Centers may review their reports at any time by accessing a password protected web page. Alternatively reports may be transmitted via e-mail or hard copy means.

Exemplary report types for monitoring produce outcomes for ambulatory surgery procedures may include, for example:

- Cumulative Reports: These reports include an accumulation of all data over a specified period of time and will therefore have a large volume/database. These reports may be used for benchmarking a center's averages with averages for all participating centers.

- Quarterly Reports: These reports include data for the previous quarter. The quarters are per calendar year, with three months in each quarter. First quarter includes January - March data; second quarter includes April - June; third quarter July - September, and fourth quarter October - December, for example. The primary purpose of the Quarterly Reports to track changes from quarter one quarter to the next.
- Current Month Reports: Monthly reports are provided to help identify individual problem cases by including only small portions of data. As a general rule, great care should be taken if using monthly reports for other purposes as the volume of data is often too small to be conclusive of any findings.
- Age Distribution is a report that is helpful when information is needed about age prevalence and/or distribution by procedure. All data submitted for each procedure group have been broken down by age, in approximate 10-year increments, for example. This report also gives the average age for that procedure.

- Recovery Time shows a breakdown in 30 minute intervals of recovery time for a specific procedure, along with the average recovery time for a center and the average time for all centers.
- 5 • Surgery Time shows a breakdown in 30 minute intervals of surgery time for a specific procedure, along with the average surgery time for a center and the average time for all centers.
- Pain, Complications and Patient Satisfaction is a  
10 table showing totals for all data specific to the quality indicators being monitored.
- General Indicators shows the rate of compliance to each indicator for a center, and the norm for "all" centers participating in the system. The  
15 general indicators report provides a view of a center's outcomes, benchmarked with other centers, at a glance.
- Complications by Payor shows a breakdown by payor classification for all complications reported.
- 20 • Complications by Anesthesia shows a breakdown is by anesthesia type (general, epidural, etc.).

Both newly generated, as well as previous reports, may be accessed by the user.

A report will typically have a defined set of benchmarking indicators. For example indicators in an exemplary embodiment of a benchmarking report for an ambulatory surgery performance outcomes monitoring may

5 include:

Perioperative Indicators

- 10
1. Patients experiencing complications of surgery during the perioperative period.
  2. Patients retained beyond the expected recovery time for the surgical procedure.
  3. Patients returned to surgery.
  4. Patients admitted to the hospital following surgery.
  - 15 5. Patients expressing pain who did not get relief of pain.

Post-Discharge Indicators

- 20
6. Patients without significant problems after discharge.
  7. Patients expressing pain after discharge who had relief of pain after utilizing pain control methods as instructed.
  8. Patients satisfied with preoperative, intraoperative, and postoperative care.
  - 25 9. Patients who received and understood discharge instructions.
  10. Patients adequately prepared for self-care at home after discharge.

30 Exemplary indicators for an embodiment of the invention for benchmarking ambulatory surgical procedures are described in detail in Example 2 below.

Referring to Figure 2, in step S800, the user reviews reports. The user, as defined herein, includes all

individuals or entities that utilize the benchmarking results. The user may receive a hard copy report or, in preferred embodiments, access the reports via a restricted web page. Each report contains specific results from the user's center (user entity) only or comparisons of the user center's performance to a norm for a group of centers. In an exemplary embodiment, the user is not allowed access to specific reports or specific results for other user centers.

In embodiments utilizing a web page to post results, it is preferable that the web page be restricted or protected such that only an authorized user for a center may access that center's customized report. This may be accomplished through use of known means such as passwords and the like.

In embodiments utilizing a web page for disseminating reports, various features may be included such as a brief definition of each indicator may be shown toward the middle of the report. Further, in an embodiment in which the report is retrieved from a web page, the user may click on the abbreviated wording for a full description of the individual indicator.

User utilization of the reports may, in an exemplary embodiment for ambulatory surgery centers, include the following:

The user may begin by assessing the volume of cases by procedure. As a general rule, at least 30 cases or more are needed before any reports can be considered as statistically significant.

5 If, for example, data on 866 knee arthroscopies were submitted for a Surgical Center from 5/17/97 - 5/31/99, this is a considerable volume for this procedure and therefore, the interpretation of reports will have statistical significance. However, if only five (5) cases were  
10 submitted for a quarter and one (1) case has complication, the rate of complications will be reflected as 20 percent. As a center continues to report more cases, this percentage may fall considerably. Thus, results based on low volumes may be useful but more antidotal in nature and should not be  
15 considered to be statistically significant.

Users have a selection of reports to choose from. In order to make it easier to go from one report to another when viewing the reports from the web page, "links" may be provided to other the benchmarking reports. These links  
20 allow the user (when in the on-line mode) to quickly switch from procedure to procedure or from report type to report type. For example, a user can get a quick look at "Surgery Time" for each of their procedures with one click or can

also assess all indicators for a given procedure by simply clicking on a link provided on the web page, for example.

An exemplary Quarterly Report showing a comparison of two data sets is provided in Figure 4. Referring to Figure 4, the circles reflect a comparison of a center's average to that of the average for all centers. A black circle indicates a center's compliance was "worse than average," a circle with a dot in the middle indicates a center's compliance was within the "average" for all centers, and a white circle indicates compliance rate was "better than the average" reported for all centers.

The arrows reflect changes in compliance rates from the previous quarter (example: fourth quarter rates are compared to third quarter rates, etc.). An arrow pointing downward indicates that the rate this quarter was "worse" than last quarter; arrows going in a straight line indicate no change from last quarter, and arrows pointing upwards indicate the rate has improved from the previous quarter.

Alternatively, data may be shown in bar graph form including graphic indication of tolerance or statistical variability, or the like. Figures 22 and 23 are exemplary of graphic forms of reports.

SOFTWARE OF THE INVENTION

The software of the invention includes software developed for the purpose of analyzing collected outcomes data and preparing reports (collectively the Report System).

5 Among the many features of the Report System software are algorithms that combine data; establish norms; compare data groups; compare data groups to norms; optionally supply statistical data; generate reports from data that is discreetly stored such that various data can be combined to  
10 generate a highly customized report in a facile manner; and archive data, combined data and reports both for referral and comparison with data acquired at a later date. The functions of the software are organized in modules. In an exemplary embodiment the modules may include a data-  
15 calculation module, data table creator module, chart generator module, comparison table module and report generator module for example. These modules are directed to the tasks of mathematically manipulating data, creating data tables, preparing charts, preparing data comparisons and  
20 generating reports, respectively. Each of these modules may have submodules directed to a specific task or type of report. Optionally, additional modules may be included to further address specialized user needs. In the preferred embodiment the reports may be accessed by a user via a



restricted web page such as a password protected web page  
for example and the web page may be suitably equipped with  
links such that additional reports, other comparisons,  
explanatory information and the like may be accessed by a  
5 simple click, for example.

A flow chart for the structure of one exemplary  
embodiment of a report system 1000 of the invention is  
provided in Figure 5.

As Figure 5 shows, the structure is subdivided into  
10 folders. The folders include among others a database folder  
1100, a configuration files folder 1200, a folder for  
storage of log files 1300, a list folder 1400 containing  
files related to mathematical calculations and layout of  
reports, a first templates folder 1500 which stores ready-  
15 to-use configuration files for different types of reports, a  
programs folder 1600 containing all modules of the report  
system and a second template folder 1700 with files for  
generating specific report features. We note that exemplary  
file contents for files of an exemplary embodiment of the  
20 invention for monitoring outcomes for ambulatory surgical  
procedures is provided in Example 1.

As one skilled in the art will recognize the exact  
composition and number of folder and files may vary  
considerably within the scope of the invention as the

invention is applied to a specific industry or service or as a user group within an industry desires customization.

However, the mathematical algorithms used and the general scheme for preparing reports are applicable to the various

5   embodiments of the invention including embodiments for benchmarking performance outcomes in ambulatory surgery, performance outcomes for other medical procedures, service performance outcomes, sales force performance outcomes, manufacturing performance outcomes and the like.

10       The report system will be described in further detail for one embodiment of the invention for monitoring performance outcomes of ambulatory surgical procedure centers.

Referring to Figure 5, a first folder is the database  
15   folder 1100. This is the main database for the report system. In the exemplary embodiment there are two tables inside this file: MEDREC and PATINT2. MEDREC, contains all medical records and PATINT2 contains all patient interview records. This file is used by the Data\_Calculator module  
20   (see the description of this module below).

Optionally the folder may also include an archive folder or backup folder or both to store incremental files.

A second folder is the configuration files folder ([INI] folder) 1200. This folder contains the configuration

files. These configuration files are designated for an end user and they allow the user to define "what reports will be generated". The files in the exemplary INI folder include Report.ini. This file is used to define the periods to  
5 generate reports for. The format of the file may be the following: Name\_of\_Parameter=Value, for example.

Comments can also be used by typing a ";" character in the beginning of the comment line. Everything after the ";" character may be ignored by the report system performing  
10 calculations and analyzing data in the exemplary embodiment. Almost all modules of report system use this file of the type.

An example of the parameters of this exemplary report file may be found in Figure 6.

15 Other configuration files may also be included. For example a file (Info file) may be included to provide paths to the different components of the report system. The info file may be used by all modules. An example format of the Info file is: Name\_of\_Parameter=Value. Exemplary Parameters  
20 for an Info file are shown in Figure 7.

Referring to Figure 5, a third folder is the Log folder 1300. This folder is used to store log files that are generated by different modules of report system. Files in this folder may include, for example, an

Executive\_Table.file which is generated by Executive\_Table  
module of the report system; an  
Executive\_Table\_Paper\_Reports, which is generated by  
Executive\_Table\_Paper\_Reports module of the report system; a  
5 DataTable.file, which is generated by DataTable module of  
the report system; a Data\_Calculator.file, which is  
generated by Data\_Calculator module of the report system; a  
GrabFile.file, which is generated by GrabFile module of the  
report system; a ProcDistrib.file, which is generated by  
10 ProcDistrib module of the report system; a report.file,  
which is generated by report module of the report system;  
and a Comparison\_Table.file, which is generated by  
Comparison\_Table module of the report system.

Referring to Figure 5 a fourth folder is a list folder  
15 1400. This folder contains configuration files like the  
[INI] folder does. But unlike the [INI] folder, almost all of  
these files are responsible either for mathematical  
calculations or for the layout of the reports. Most of  
these files are a list of strings that have the same  
20 structure. Usually each string consists of several fields  
and the fields are separated by "\*" character.

Files in the list folder may include for example an  
Executive\_Table list file, which is used by Executive\_Table  
module. This list file contains a list of indicators that

will be shown in a preferred embodiment of a report. The format of each string may be the following:

Indicator\_Name\*Numerator\*Denominator\*Description\*MinOrMax\*Link

5 Exemplary Executive\_table contents are shown in Figure 8.

An example of parameters associated with this list file is shown in Figure 4.

Another list file may be a DataTableItems list file.  
10 The DataTableItems list file is used by DataTable\_Creator module. The DataTable\_Creator module generates "Data Tables" file ("DataTable.html"). DataTableItems file may consist of several sections. Each section corresponds to a separate table in "DataTable.html" file. Strings that are  
15 located inside each section are used to customize the rows in tables. Each section may begin with one of the following strings:

---Name\_of\_Table\*Total\_by\_Proc\_Flag or  
Name\_from\_Dump\_DB\*Row\_Name

20 Figures 9 and 10 show examples of the data table items file parameters and associated data tables, respectively.

Another list file may be an Indicators file which is used by the Chart\_Generator module. The Indicators file consists of several sections. Each section corresponds to a

5 section may begin with the following string:

With\_Tolerance\*Without\_Tolerance\*Chart Header\*Chart Footer

10           Indicators for each section are described after the  
section header. An example format of indicator description  
string is the following:

Figures 12 and 13 show examples of an indicator  
15 description string and report page, respectively.

20        Message\_Name=Text\_of\_Message

      where Message\_Name is a name of message (report system modules use this name to refer to log messages) and Text\_of\_Message is a message text. Usually a message text is much longer than Message\_Name, so the main purpose of LogMessages list file is to eliminate extra text and also

this list file allows similarity of log messages throughout all modules.

Another list file may be a Comparison\_table list file which is used by the Comparison\_table module. It is divided into several sections. Each section begins with a header and represents a separate group of indicators in "Comparison Table". (This may be used with paper reports, for example). The format of the header is:

---Header\_of\_indicator\_group\*

wherein Header\_of\_indicator\_group contains a text that will be used as a header of the indicator group. If Header\_of\_indicator\_group is empty then no header is used and all indicators in this group have a bold font, for example.

Each section is followed by indicator definitions. The format of these definitions may be the following:

Numerator\*Denominator\*Descripting\_Text\*

An example of the parameters of a Comparison table file is found in Figure 14.

A Corporate\_Members list file may be included also. In an exemplary embodiment, this file describes corporate members or unit entities using the benchmarking system. Each unit entity is represented by a separate section. All sections may have a header. The format of headers may be:

---GroupName\*GroupUsername\*GroupUserCode\*members\_access

where GroupName may be a name of the unit entity;

GroupUsername may be a username of the group (this username may be used to access reports); GroupUserCode may be a

5 usercode of the group(a three letter code may be convenient in some embodiments); members\_access ("Yes" or "No") may be used to restrict the access of separate members (in individual users) to the reports. members\_access may be used by "New\_Center\_Prepare" module to customize Apache  
10 ".htaccess" files for example. Separate members (individual users) of the user entity can access their reports if members\_access="Yes" otherwise only GroupUsername can be used to access the reports.

The header of the section may be followed by a list of  
15 usercodes of centers that belong to the group. The usercodes may be separated by "Enter" key.

An example of a report of the type which may be presented to a user showing only data for that user and cumulative data for all user groups included in establishing  
20 the norm is shown in Figure 15. Thus, access to a unit entity's individual results may be restricted to that unit entity.

Another file may be a ProcConv list file. This file is used by Appender and Data\_Calculator. This file defines a



mapping table between the CPT codes and the procedure groups. When Appender module appends new data it ignores the "PROC" fields in incremental files and uses this mapping, in the same way Data\_Calculator ignores the existing "PROC" field in the report master database and recreates this field using the mapping. All strings in ProcConv.lst file may have the same format such as:

CPT\_Code\* Procedure\_Group

where CPT\_Code is a five digit CPT code (procedure code) and Procedure\_Group is the name of the corresponding procedure group.

Another list file may be a ProcDistrib file which is used by ProcDistrib module. It defines what fields from "\*\_dump.mdb" databases should be displayed in the "Case Distribution" table. The format of the file may be the following:

Field\_Name\*Descripting\_Text\*

Examples of the file parameter information of a ProcDistrib and a case distribution table are found in Figures 16 and 17, respectively. Note that in the example shown in Figure 17 only two strings are shown TotMR\*Medical Records\* and TotPI\*Patient Interview\* this list can be easily expanded.

Another list file may be Sites list file or its equivalent which contains information about centers - it describes relationship between usercodes, real names and usernames. The format of this file may be:

5    Three\_Letter\_USERCODE\*real\_Name\_of\_center\*Centers\_Username  
where Three\_Letter\_USERCODE is the usercode of the center;  
real\_Name\_of\_center is the real name of the center and  
Centers\_Username is the username. Note that string ALL\*All  
centers\* MUST be first in some embodiments as this string  
10 defines usercode for all centers.

Another list file of the exemplary embodiment is a first Stage list file, (Stage1.lst) which is used by Data\_Calculator module only and describes the mathematical expressions using the MS SQL language. To get a table that  
15 contains procedure level data for a certain period,  
Data\_Calculator module runs in two stages. On the first stage, the module uses a first stage list file (Stage1.lst) to create a "SELECT"-query that combines MEDREC and PATINT2 tables in one table, for example. Instead of the original  
20 fields, this table contains new calculated fields that are used to calculate fields in "\*\_dump.mdb" files on the next stage. In this stage, the records are not grouped by procedure groups and centers usercodes - they are still patient-level records.

0996475.12001

The format of strings of the first stage list file may be the following:

Name\_of\_Field\_1\*SQL\_Expression

Alternatively a simplified version of "SELECT"-query  
5 for use in the first stage may be written in the following way:

SELECT expression\_1 AS field\_1,...,expression\_k AS  
field\_k,...,expression\_N AS field\_N FROM table

10 Name\_of\_Field is used as *field\_k* and SQL\_Expression is used as *expression\_k*.

A second Stage list file (Stage2.lst), which is also used by the Data\_Calculator module, describes mathematical  
15 expressions using MS SQL language. This file is used on the second stage of the calculation of MasterTable (later, records from MasterTable are used to populate "\*\_dump.mdb" files). On this stage MasterTable is calculated. It contains procedure-level data. Only this procedure-level data is used  
20 by other modules on next the steps of the report generation process.

The format of strings of this file may be the following:

Name\_of\_Field\_2\*Data\_Type\_of\_Field\*SQL\_Expression\*Denominator\*  
25

Exemplary Stage files are shown in Figures 18 and 19.

Referring to Figure 5, a fifth folder is the first  
Templates folder 1500. Files in this folder are not used  
directly by the report system. The purpose of the first  
templates folder 1500 is to store ready-to-use configuration  
5 files for different types of reports. For example, one can  
use this folder to store configuration files used to  
generate sample reports. As shown, this folder contains two  
files only (Sample-Sites.lst and Full\_List\_Sites.lst) - they  
are two versions of Sites.lst file: one is for the usual  
10 reports, another for sample reports.

Referring again to Figure 5, a sixth folder is the  
Programs folder 1600. This folder contains all report  
modules of the report system. As shown, all modules may be  
divided into two groups: for "Paper Quarterly Reports" and  
15 for "Web Reports". [Paper\_Reports] and [Web\_Reports]  
folders were created according to this breakdown.

The [Paper\_Reports] subfolder contains modules that are  
used to generate paper reports. For example, for the "Paper  
Quarterly Report," HTML files generated by the paper reports  
20 modules are created in the folder defined by  
"SavePathForPaperReport." An exemplary structure of the  
"Paper\_Reports" folder is:

```
25      [PAPER_REPORTS]
      +---[Comparison_Table]
      |   |
```

```

    |      +---[Qyyyyy-qq]
    |      |
    |      +---XYZ-FullNameOfXYZ.html
5   |
    | +---[Executive_Table]
    | |
    | +---[Qyyyyy-qq]

```

Another exemplary report module is the executive table  
10 (Executive\_Table\_Paper\_Reports.exe). This module generates  
"Executive table" for a paper quarterly report. It may  
create HTML files for all quarters defined in a report.ini  
file.

An example of an executive report is shown in Figure 4.

15 Another exemplary report module (Comparison\_table.exe)  
generates the "Comparison table" for the paper quarterly  
report. It may create HTML files for all quarters defined  
in the "report.ini" file.

A Web\_Reports subfolder or its equivalent contains  
20 modules of the report system that generate reports to be  
provided to the user through the Internet or network.

A first file APPENDER or its equivalent may appends  
incremental files into "Report Master Database" file. It  
may use only one configuration file. This module may scan  
25 all centers directories located under upload directory  
defined by parameter "UploadDirectory" and append  
incremental records from these folder to the master

database. Appended incremental files are moved to the  
centers "Backup" folder located under their upload  
directories.

A second file Chart\_Generator or its equivalent may  
5 generate the large part of reports for internet  
dissemination. Chart Generator may, for example, create the  
following reports in one embodiment:

- 10
1. Age Distribution,
  2. Recovery Time,
  3. Surgery Time,
  4. General Indicators,
  5. Complications by Payor,
  6. Complication by Anesthesia.

15 This module may use input files including:

- 20
1. Corporate\_Members.lst
  2. Indicators.lst
  3. "\*\_dump.mdb" files
  4. MasterTable.mdb
  5. Sites.lst
  6. LogMessages.lst
  7. all HTML template files from folder, defined by  
"TemplateDirectory" parameter in "new-soix.ini"

Reports may be divided into two types: the reports  
25 which contain data for all procedure groups in one page and  
those that have separate page for each procedure group.  
Report folders for all periods have special sub-folders for  
each procedure group (in Fig. 20 these folders are shown as  
[Proc\_1], [Proc\_2], [Proc\_k] and [Proc\_Z]) and there are the  
30 index files (in Fig. 20 these files are referred as

report\_index\_file\_1.html, . . . ., report\_index\_file\_M.html)  
that contain links to this report pages. All these folders  
and files are generated by the Chart\_Generator. Also, this  
module may refresh main-Old.html, the main page of each  
5 center (index.html file that are located directly in the  
center's folder, not in sub-folders), and index.html files  
for all recalculated periods.

If a center has a custom picture that is included in  
"Sites.lst" then its main page and main picture will be  
10 updated by Chart\_Generator.exe.

In one exemplary embodiment the Chart\_Generator.exe  
module differs from other report modules. When other report  
modules are running they update only the report files that  
they generate - they do not delete any other report files,  
15 so there is no need to rerun other modules later. When the  
Chart\_Generator module is running, it may delete the whole  
report folder for a given period.

Other exemplary report modules include an  
Executive\_Table.exe module which creates "The Executive  
20 Benchmark Table" tables for web page quarter reports. As  
input files it uses:

1. Sites.lst
2. Corporate\_Members.lst
3. Executive\_Table.lst
- 25 4. LogMessages.lst
5. "\*\_dump.mdb" files

6. report.ini
7. new-soix.ini

As output files, it creates "Executive\_Table.html"

5 files for all quarters defined in "report.ini" file. This module can be easily modified to generate the reports for other periods, in addition to quarters. An exemplary executive table report is shown in Figure 4.

Another exemplary report module is the

10 DataTable\_Creator module which generates indicator results tables such as "Pain, Complication & Patient Satisfaction" tables.

A further exemplary report module may be a Data\_Calculator module as its equivalent. This module

15 calculates procedure-level data for all centers. In some exemplary embodiments only this module and "APPENDER" have direct access to the main database, all other modules just use procedure-level data. This serves to eliminate extra calculations, for example if the layout of several reports

20 is changed then there is no need to recalculate the data, one just runs the necessary modules and reports are refreshed. In most cases it takes much less time then when the data should be recalculated.

Another report module may be ProcDistrib\_Creator or its

25 equivalent. This module creates "Case Distribution" tables.



These tables are created for the whole network. They show case distribution by procedure group and by site inside each procedure group. The number of medical records and patient interviews may be present on these tables in an exemplary  
5 embodiment.

Referring to Figure 5, a seventh folder in one exemplary embodiment is a second Template folder 1700. This folder may contain, for example in an exemplary embodiment, templates for reports such as the "Age Distribution" report  
10 used by "Chart\_Generator" module; the "Complication by Anesthesia" with tolerance zone report which is used by "Chart\_Generator" module; the "Complication by Anesthesia" without tolerance zone report which is used by "Chart\_Generator" module; the "General Indicators" with  
15 tolerance zone report which is used by "Chart\_Generator" module; the "General Indicators" without tolerance zone report which is used by "Chart\_Generator" module; the "Reports for Previous Periods" which is used by Chart\_Generator module; the loopback file which is shown  
20 instead of reports when a center does not have data for certain period and is used by Chart\_Generator, DataTable\_Creator and ProcDistrib\_Creator modules; a main-old file which is used as main page for reports for previous period and contains links to these reports and used by

Chart\_Generator module; a main-Template file which is used  
as main index page for "Current Month Reports", "Quarterly  
Reports" and "Cumulative reports" by the Chart\_Generator  
module; a main.html file which is used as center's main  
5 page, includes either default SOIX picture or a favorite  
picture of the center and is used by Chart\_Generator module;  
a Payor.html file which is a template for "Complications by  
Payor" reports with tolerance limits and is used by  
Chart\_Generator module; a Payor2.html file which is a  
10 template for "Complications by Payor" reports without  
tolerance limits and used by Chart\_Generator module; a  
RecovTime2.html file which is a template for "Recovery Time"  
reports and is used by Chart\_Generator module; and a  
SurgTime2.html file which is a template for " Surgery Time"  
15 reports and is used by Chart\_Generator module.

A subfolder in the template 1700 folder of an exemplary  
embodiment is an All folder. This folder contains files  
that are used for reports for the whole network and has the  
same application as corresponding templates in the first  
20 template folder 1500.

Another exemplary subfolder of the templates 1700  
folder is an image folder which stores various images that  
are used for reports. The image folder may contain a  
centers folder. This folder may be used to store a center's

favorite pictures in a format such as "USERCODE.(jpg or gif)", where USERCODE is three-letter user code of a center (For example: aaa.jpg, aza.gif).

In a preferred embodiment a user may access the reports prepared via a web page. This web page may be password protected to restrict user access or selectively release specific data to a specified user or both. An exemplary report folder structure for a private web site is shown in Figure 20.

For this example, folders [center\_1], ..., [center\_k], ..., [center\_Y] are the centers report directories corresponding to each center. The name of a center's directory may be a three letter USERCODE of the center for example: AAA, AAB, MFA. Each center has its own home page, for example [center\_k]\index.html file.

Each center's folder has file structure which is described below for an exemplary embodiment.

Folders [Month], [Quarter] and [MonthCumul] are used to store "Current Month Report", "Quarterly Reports" and "Cumulative Reports", respectively.

[OLD] folder is used to store "Reports for the Previous Periods". Each folder inside [OLD] represents reports for a certain period. Folders for quarterly reports have [Qyyyy-qq] names, where "yyyy" is a year in four digit format and

"qq" is a quarter number with a leading zero (Example:  
"Q1999-03" - reports for third quarter of 1999). Files for  
previous monthly reports are located in [Myyyy-mm] folders,  
and cumulative monthly reports are saved in [Myyyy-mmC]  
ones, where "yyyy" is a year and "mm" is a month (Example:  
"M1999-04" - monthly reports for April 1999, "M1999-03C" -  
cumulative monthly reports for March 1999). A file shown in  
Figure 20 as main-Old.html provides links to reports for the  
previous periods. An example of a menu that permits a user  
to use this file is shown in Fig. 21.

Referring again to Figure 20, a folder such as Download  
may be used to store center specific files, like  
"ftprun.run" or updates and patches for OMS program.

Other Database files such as (Month\_dump.mdb,  
MonthCumul\_dump.mdb and Quarter\_dump.mdb) may contain  
procedure-level data calculated by the report system. These  
files store data for "Current Month Reports", "Cumulative  
Reports" and "Quarterly Reports" respectively. In an  
exemplary embodiment almost all parts of the report system  
use these files - not the original Medical Record and  
Patient Interview patient-level tables.

All report pages may be divided into the two groups:

- 1) report pages that show information only for a  
specific procedure group (in current version of

the report system these files include a chart and a table corresponding to this chart);

- 2) report pages that include information for all procedure groups on one page (in current version of the report system these files include tables only).

Index files which do not actually include any reports, may be used to provide access to report pages. Index files may include the following in an exemplary embodiment, for example: an Index file for "Age Distribution" reports; an Index file for "Complication by Anesthesia" reports; an Index file for "General Indicators" reports; an Index file for "Complications by Payor" reports; an Index file for "Recovery Time" reports; and an Index file for "Surgery Time" reports.

Combined files are report files for reports where data for all procedure groups are combined in one file. These files may contain tables only. Examples of such files may include "The Executive Benchmark Table" report table or "Data Tables" report tables, for example.

Another file may be included which is used by the report system to determine the period of reports.

Folders for each procedure group may be included for preparing reports having separate pages for each procedure

5 all spaces to "\_" (For example: the folder for procedure  
group "D&C/Hysteroscopy" is "DCHysteroscopy", "ENT - T&A <  
12" is "ENT\_-\_TA\_\_12"). Each report in these exemplary  
folders is represented by two files: *report\_file\_k.html* and  
*report\_file\_k.gif*. The HTML file includes a table and a  
10 chart in GIF format (For example: "Age Distribution" for  
procedure group "Carpal Tunnel" is represented by the files  
"Age\_Distribution2.html" and "Age\_Distribution2.gif" in the  
folder "CarpalTunnel").

```
15  the following order to create a report:  Data_Calculator;
    Chart_Generator; Executive_Table; DataTable_Creator; and
    ProcDistrib Creator.
```

The order of each module should be completed before moving to the next.

20        Software will also contain suitable provision for  
adding and removing user entitles from the benchmarking  
system.

EXAMPLE 1

The following is a listing of exemplary files contents  
for data analysis and report preparation for one embodiment  
of the invention. This example exemplifies one of many  
5 groups of files which may be used in ambulatory surgical  
procedures monitoring and is provided as an illustrative  
example of the many configuration file types and components  
which are within the scope of the invention.

**Report.ini** file

10 ;Everything that is located after ";" is comments  
  
[Run options]  
CalculationDate = ;1/30/2000 ;Date that will be used as creation date in  
charts and tables  
15 ;If there is no date then current date is  
used  
  
MinNumberOfCases =20 ;Minimal number of cases that allows to  
generate reports for the procedure group  
20 Confidence =1.7 ;Coefficient before sigma to calculate  
tolerable limits  
  
FoundationDate =1/1/1999  
  
25 QuarterlyReports =yes ;Yes or No  
QuarterStart =1 ;1,2,3 or 4  
QuarterYearStart =2000 ;format is yyyy  
QuarterEnd =2 ;1,2,3 or 4  
QuarterYearEnd =2000 ;format is yyyy  
30 MonthlyReports =yes ;Yes or No  
MonthStart =1 ;1 to 12  
MonthYearStart =2000 ;format is yyyy  
MonthEnd =6 ;1 to 12  
35 MonthYearEnd =2000 ;format is yyyy  
  
CumulativeMonthlyReports =yes  
CumulativeMonthStart =1  
CumulativeYearStart =2000  
40 CumulativeMonthEnd =6  
CumulativeYearEnd =2000  
  
;-----parameters below this line are not supportes at this moment  
45 RunMode =Auto ;Auto or Manual  
  
CleanedDBF =no ;Yes or No

StandardReport =no ;Yes or No  
StartDate =1/1/1998  
EndDate =12/31/1998

5

*Executive\_Table.lst* file

ind1\*Nummer1\*TotMR\*Perioperative Compli-<br>cations\*min\*/genrep/ind1.htm  
ind2\*Nummer2\*TotMR\*Delayed in Discharge\*min\*/genrep/ind2.htm  
ind3\*Nummer3\*TotMR>Returns to Surgery\*min\*/genrep/ind3.htm  
10 ind4\*Nummer4\*TotMR\*Admits to Hospital\*min\*/genrep/ind4.htm  
ind5\*Nummer5\*Denom5\*Pain Episodes Not Relieved<br>\*min\*/genrep/ind5.htm  
  
IND6\*Nummer6\*TotPI\*Care Not Needed After<br> Discharge\*max\*/genrep/ind6.htm  
IND7\*Nummer7\*Denom7\*Pain Controlled After<br> Discharge\*max\*/genrep/ind7.htm  
15 IND8\*Nummer8\*TotPI\*Satisfied Patients\*max\*/genrep/ind8.htm  
IND9\*Nummer9\*TotPI\*Effective Discharge <br>Instructions\*max\*/genrep/ind9.htm  
IND10\*Nummer10\*TotPI\*Patients Prepared for <br>Self-Care\*max\*/genrep/ind10.htm

20 *DataTableItems.lst* file

---PATIENT DISPOSITION\*TotByProc  
Patient\_Dispos\_Normal\*Normal  
Patient\_Dispos\_RetainedMore3Hrs\*Retained >3 hrs  
Patient\_Dispos\_Hospital\*Hospital  
25 Patient\_Dispos\_Reoperated\*Reoperated  
  
---ANESTHESIA\*TotByProc  
TotEpi\*Epidural .  
TotGen\*General  
30 TotSpi\*Spinal  
TotMAC\*MAC  
TotBlock\*Block  
TotTopical\*Topical  
TotLoc\*Local  
35 TotIVC\*IV-CON SED  
TotOther\*Other  
TotNone\*None  
  
---PAIN AND COMPLICATIONS\*  
40 Pain\_Complic\_NoPain-NoComplic\*No Pain, No<br>Complications  
Pain\_Complic\_Pain\*Pain  
Pain\_Complic\_Nausea\*Nausea  
Pain\_Complic\_Vomiting\*Vomiting  
Pain\_Complic\_InabilityToVoid\*Inability To Void  
45 Pain\_Complic\_Bleeding\*Bleeding  
Pain\_Complic\_InstabVitalSigns\*Instability Of<br>Vital Signs  
Pain\_Complic\_LevelOfConscChanges\*Level Of Conscious-<br>ness  
Changes  
Pain\_Complic\_RespirProblems\*Respiratory<br>Problems



0996475.12001  
T002T 52495550

---PAIN CONTROL METHODS\*  
Pain\_Control\_Meth\_PainContrMethExplOnDischarge\*Pain  
Control<br>Methods Explained<br>on Discharge  
5 Pain\_Control\_Meth\_PrescrGivenOnDischarge\*Prescription<br>Given  
On Discharge  
Pain\_Control\_Meth\_PainVerb\*Pain Verbalized  
Pain\_Control\_Meth\_MedOrdered\*Medication<br>Ordered<br>(Who Had  
Pain)  
10 Pain\_Control\_Meth\_MedAdmin\*Medication<br>Administered<br>(Who  
Had Pain)  
Pain\_Control\_Meth\_MedAdminAndRefused\*Medication<br>Administere  
d<br>And Refused<br>(Who Had Pain)  
Pain\_Control\_Meth\_PainRelieved\*Pain Relieved<br>(Who Had Pain)  
15 ---AFTER LEAVING THE SURGERY CENTER\*  
After\_Leave\_Surgery\_Problems\_Might\_Have\*Knew What  
Problrms<br>Might Have  
After\_Leave\_Surgery\_Who\_Call\*Knew Who To Call  
20 After\_Leave\_Surgery\_Meds\_To\_Use\*Knew What<br>Medicines to Use  
After\_Leave\_Surgery\_Had\_Appointment\*Had an Appointment  
After\_Leave\_Surgery\_Had\_All\_Info\*Had All Information  
  
---POSTOPERATIVE PATIENT INTERVIEW:<br>COMPLICATIONS THAT  
25 REQUIRED MEDICAL INTERVENTION\*  
Postop\_Pat\_Int\_Complic\_AnyProblem\*Any Problem  
Postop\_Pat\_Int\_Complic\_Nausea\*Nausea  
Postop\_Pat\_Int\_Complic\_Vomiting\*Vomiting  
Postop\_Pat\_Int\_Complic\_Fever\*Fever  
30 Postop\_Pat\_Int\_Complic\_ProblemUrine\*Problem Urinating  
Postop\_Pat\_Int\_Complic\_Bleeding\*Bleeding  
Postop\_Pat\_Int\_Complic\_SignsOfInf\*Signs Of Infection  
  
---PAIN MANAGEMENT AT HOME\*  
35 Pain\_Manag\_Home\_PostopPainAtHome\*Postop Pain<br>at Home  
Pain\_Manag\_Home\_PostopInstrContrPain\*Postop Instructed<br>to  
Control Pain<br>at Home  
Pain\_Manag\_Home\_ComplWithInstr\*Complied with<br>Instructions  
  
40 ---PAIN RELIEF AT HOME FOR PATIENTS WHO HAD PAIN\*TotByProc  
Pain\_Relief\_Home\_Completely\*Completely  
Pain\_Relief\_Home\_Greatly\*Greatly  
Pain\_Relief\_Home\_Somewhat\*Somewhat  
Pain\_Relief\_Home\_NotRelieved\*Not Relieved  
45 ---PERCEIVED QUALITY IN REGISTRATION AND ADMISSION  
PROCESS\*TotByProc

Perceived\_Quality\_Reg\_And\_Admis\_Excellent\*Excellent  
Perceived\_Quality\_Reg\_And\_Admis\_Good\*Good  
Perceived\_Quality\_Reg\_And\_Admis\_Fair\*Fair  
Perceived\_Quality\_Reg\_And\_Admis\_Poor\*Poor  
5 Perceived\_Quality\_Reg\_And\_Admis\_N-A\*N/A

---PERCEIVED QUALITY AT PREADMISSION TESTING\*TotByProc  
Perceived\_Quality\_Preadmis\_Excellent\*Excellent  
Perceived\_Quality\_Preadmis\_Good\*Good  
10 Perceived\_Quality\_Preadmis\_Fair\*Fair  
Perceived\_Quality\_Preadmis\_Poor\*Poor  
Perceived\_Quality\_Preadmis\_N-A\*N/A

---PERCEIVED QUALITY IN RECOVERY STAGE IN THE CENTER\*TotByProc  
15 Perceived\_Quality\_Rec\_Stage\_Excellent\*Excellent  
Perceived\_Quality\_Rec\_Stage\_Good\*Good  
Perceived\_Quality\_Rec\_Stage\_Fair\*Fair  
Perceived\_Quality\_Rec\_Stage\_Poor\*Poor  
Perceived\_Quality\_Rec\_Stage\_N-A\*N/A  
20

*Indicators.lst* file

;General Indicators  
---ind\*ind2\*General Indicators\*%Average% Records:  
25 %TotMRAllSites%, %You%: %TotMRThisSite%. %Average%  
Interviews: %TotPIAllSites%, %You%: %TotPIThisSite%.  
ind1\*Nummer1\*TotMR\*ind1  
ind2\*Nummer2\*TotMR\*ind2  
ind3\*Nummer3\*TotMR\*ind3  
30 ind4\*Nummer4\*TotMR\*ind4  
ind5\*Nummer5\*Denom5\*ind5^^  
IND6\*Nummer6\*TotPI\*IND6  
IND7\*Nummer7\*Denom7\*IND7^^  
IND8\*Nummer8\*TotPI\*IND8  
35 IND9\*Nummer9\*TotPI\*IND9  
IND10\*Nummer10\*TotPI\*IND10

;ind1 by payor  
---payor\*payor2\*Complications by Payor\*%Average% Average  
40 Complication Rate: %Ind1AllSites%%. %You%: %Ind1ThisSite%%.  
Care\_Ind1\*Care\*TotCare\*Medicare  
Aid\_Ind1\*Aid\*TotAid\*Medicaid  
Com\_Ind1\*Com\*TotCom\*Non-Capitated  
Cap\_Ind1\*Cap\*TotCap\*Capitated  
45 Uni\_Ind1\*Uni\*TotUni\*Uninsured  
Wor\_Ind1\*Wor\*TotWor\*Workmens Comp

Oth\_Ind1\*Oth\*TotOth\*Other

;ind1 by anesthesia

---anest\*anest2\*Complications by Anesthesia\*%Average%

5 Average Complication Rate: %Ind1AllSites%. %You%:  
%Ind1ThisSite%.

EPI\_Ind1\*EPI\*TotEPI\*Epidural

GEN\_Ind1\*GEN\*TotGEN\*General

Spi\_Ind1\*Spi\*TotSpi\*Spinal

10 MAC\_Ind1\*MAC\*TotMAC\*MAC

Block\_Ind1\*Block\*TotBlock\*Block

Topical\_Ind1\*Topical\*TotTopical\*Topical

Loc\_Ind1\*Loc\*TotLoc\*Local

IVC\_Ind1\*IVC\*TotIVC\*IV-CON SED

15 Other\_Ind1\*Other\*TotOther\*Other

None\_Ind1\*None\*TotNone\*None

;Surgery Time

---\*surgttime2\*Surgery Time\*Surgery Time (min)||%Average%

20 average: %Surgtime\_AvgAllSites% min. %You%:

%Surgtime\_AvgThisSite% min.

SURGTIME-0-30V\*SURGTIME-0-30\*SURGTIME\_TOT\*0-29

SURGTIME-30-60V\*SURGTIME-30-60\*SURGTIME\_TOT\*30-59

SURGTIME-60-90V\*SURGTIME-60-90\*SURGTIME\_TOT\*60-89

25 SURGTIME-90-120V\*SURGTIME-90-120\*SURGTIME\_TOT\*90-119

SURGTIME-120-150V\*SURGTIME-120-150\*SURGTIME\_TOT\*120-149

SURGTIME-150-180V\*SURGTIME-150-180\*SURGTIME\_TOT\*150-179

SURGTIME-180-210V\*SURGTIME-180-210\*SURGTIME\_TOT\*180-209

SURGTIME-210-240V\*SURGTIME-210-240\*SURGTIME\_TOT\*210-239

30 SURGTIME-240+V\*SURGTIME-240+\*SURGTIME\_TOT\*240+

;Recovery Time

---\*recovtime2\*Recovery Time\*Recovery Time (min)||%Average%

average: %Rectime\_AvgAllSites% min. %You%:

35 %Rectime\_AvgThisSite% min.

RECTIME-0-30V\*RECTIME-0-30\*RECTIME\_TOT\*0-29

RECTIME-30-60V\*RECTIME-30-60\*RECTIME\_TOT\*30-59

RECTIME-60-90V\*RECTIME-60-90\*RECTIME\_TOT\*60-89

RECTIME-90-120V\*RECTIME-90-120\*RECTIME\_TOT\*90-119

40 RECTIME-120-150V\*RECTIME-120-150\*RECTIME\_TOT\*120-149

RECTIME-150-180V\*RECTIME-150-180\*RECTIME\_TOT\*150-179

RECTIME-180-210V\*RECTIME-180-210\*RECTIME\_TOT\*180-209

RECTIME-210-240V\*RECTIME-210-240\*RECTIME\_TOT\*210-239

RECTIME-240+V\*RECTIME-240+\*RECTIME\_TOT\*240+

45

;Age Distribution

Age\_Distrib\_0-14V\*Age\_Distrib\_0-14\*Age\_Distrib\_Tot\*0-14  
Age\_Distrib\_15-24V\*Age\_Distrib\_15-24\*Age\_Distrib\_Tot\*15-24  
Age\_Distrib\_25-34V\*Age\_Distrib\_25-34\*Age\_Distrib\_Tot\*25-34  
Age\_Distrib\_35-44V\*Age\_Distrib\_35-44\*Age\_Distrib\_Tot\*35-44  
Age\_Distrib\_45-54V\*Age\_Distrib\_45-54\*Age\_Distrib\_Tot\*45-54  
Age\_Distrib\_55-64V\*Age\_Distrib\_55-64\*Age\_Distrib\_Tot\*55-64  
Age\_Distrib\_65-74V\*Age\_Distrib\_65-74\*Age\_Distrib\_Tot\*65-74  
Age\_Distrib\_75-84V\*Age\_Distrib\_75-84\*Age\_Distrib\_Tot\*75-84  
Age\_Distrib\_85+V\*Age\_Distrib\_85+\*Age\_Distrib\_Tot\*85+

*LogMessages.lst* file

```
KillFileMsg      = File was overwritten or deleted
```

Comparison table.lst file

68

Patent Application  
Attorney Docket No.: 58367.000003

	Pain_Control_Meth_PainContrMethExplOnDischarge*TotMR*		Percent Pain Control Methods Explained*
5	---After Leaving the Surgery Center* After_Leave_Surgery_Problems_Might_Have* TotPI* Problems They Might Have* After_Leave_Surgery_Who_Call* TotPI* After_Leave_Surgery_Meds_To_Use* TotPI* Control Pain*		Percent That Knew What Percent Knew Who to Call* Percent Knew Medications to
10	After_Leave_Surgery_Had_Appointment* TotPI* Appointment* After_Leave_Surgery_Had_All_Info* TotPI* Info*		Percent with Post Operative Percent Who Had Self Care
15	---Problems at Home* Postop_Pat_Int_Complic_AnyProblem* TotPI* to Procedure*		Percent with Problem Related
20	Postop_Pat_Int_Complic_Nausea* TotPI* Postop_Pat_Int_Complic_Vomiting* TotPI* Postop_Pat_Int_Complic_Fever* TotPI* Postop_Pat_Int_Complic_ProblemUrine* TotPI* Urinating*		&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;& Nausea* &nbsp;&nbsp;&nbsp;&nbsp;& Vomiting* &nbsp;&nbsp;&nbsp;&nbsp;& Fever* &nbsp;&nbsp;&nbsp;&nbsp;& Difficulty
25	Postop_Pat_Int_Complic_Bleeding* TotPI* Postop_Pat_Int_Complic_SignsOfInf* TotPI* Infection*		&nbsp;&nbsp;&nbsp;&nbsp;& Bleeding* &nbsp;&nbsp;&nbsp;&nbsp;& Signs of
30	Pain_Manag_Home_PostopPainAtHome* TotPI* Pain_Manag_Home_PostopInstrContrPain* Pain_Manag_Home_PostopPainAtHome* Instruction about Pain* Pain_Manag_Home_ComplWithInstr* Pain_Manag_Home_PostopPainAtHome* Instructions* Pain_Relief_Home_Completely* Pain_Manag_Home_PostopPainAtHome* Completely Relieved*		Percent Bothered by Pain* &nbsp;&nbsp;&nbsp;&nbsp;& Percent with &nbsp;&nbsp;&nbsp;&nbsp;& Percent Following &nbsp;&nbsp;&nbsp;&nbsp;& Percent
35	---Perceived Quality of Care* Nummer8* TotPI* Perceived_Quality_Reg_And_Adms_Excellent* TotPI* Registration and Admission* Perceived_Quality_Preadmis_Excellent* TotPI* Preadmission Testing*		Percent Excellent Quality* Percent Excellent Percent Excellent
40	Perceived_Quality_Rec_Stage_Excellent* TotPI* Stage*		Percent Excellent Recovery

**ProcConv.lst file**

45	29888* Arthroscopic ACL Repair 67916* Blephroplasty 67921* Blephroplasty 19325* Breast augmentation 19120* Breast Biopsy
50	19318* Breast reduction 31622* Bronchoscopy 31625* Bronchoscopy 28290* Bunionectomy 28292* Bunionectomy
55	28293* Bunionectomy 28294* Bunionectomy 28296* Bunionectomy 28297* Bunionectomy 28298* Bunionectomy
60	28299* Bunionectomy 29848* Carpal Tunnel 64721* Carpal Tunnel 66830* Cataract removal 66840* Cataract removal

66850\* Cataract removal  
66852\* Cataract removal  
66920\* Cataract removal  
66930\* Cataract removal  
5 66940\* Cataract removal  
66983\* Cataract removal  
66984\* Cataract removal  
45378\* Colonoscopy, diagnostic  
45380\* Colonoscopy with biopsy  
10 45384\* Colonoscopy with biopsy  
45385\* Colonoscopy with biopsy  
52000\* Cystoscopy  
52005\* Cystoscopy  
52007\* Cystoscopy  
15 52204\* Cystoscopy  
52281\* Cystoscopy  
58120\* D&C/Hysteroscopy  
58558\* D&C/Hysteroscopy  
43235\* EGD  
20 43239\* EGD with biopsy  
43248\* EGD with dilation  
43249\* EGD with dilation  
30520\* ENT- Septoplasty  
31255\* ENT Sinus endoscopy  
25 42820\* ENT- T&A < 12  
42826\* ENT- Tonsillectomy > 12  
69436\* ENT- Tubes  
69631\* ENT- Tympanoplasty  
49320\* GYN laparoscopy  
30 58660\* GYN laparoscopy  
58670\* GYN laparoscopy  
58671\* GYN laparoscopy  
49505\* Hernia repair  
49585\* Hernia repair  
35 29870\* Knee Arthroscopy  
29877\* Knee Arthroscopy  
29881\* Knee Arthroscopy  
29882\* Knee Arthroscopy  
29884\* Knee Arthroscopy  
40 47562\* Laparoscopic cholecystectomy  
47564\* Laparoscopic cholecystectomy  
19125\* Needle localization breast biopsy  
62310\* Pain management -epidural  
62311\* Pain management -epidural  
45 64510\* Pain management -epidural  
20550\* Pain management -injection  
55700\* Prostate biopsy

30400\* Rhinoplasty  
15828\* Rhytidectomy  
23412\* Shoulder Arthroplasty (open)  
23450\* Shoulder Arthroplasty (open)  
5 23455\* Shoulder Arthroplasty (open)  
29815\* Shoulder Arthroscopy, dx or tx  
29819\* Shoulder Arthroscopy, dx or tx  
29820\* Shoulder Arthroscopy, dx or tx  
29821\* Shoulder Arthroscopy, dx or tx  
10 29822\* Shoulder Arthroscopy, dx or tx  
29823\* Shoulder Arthroscopy, dx or tx  
29825\* Shoulder Arthroscopy, dx or tx  
29826\* Shoulder Arthroscopy, dx or tx

15

56340\* Laparoscopic cholecystectomy  
56342\* Laparoscopic cholecystectomy  
56300\* GYN laparoscopy  
20 56302\* GYN laparoscopy  
56304\* GYN laparoscopy  
56351\* D&C/Hysteroscopy  
62275\* Pain management -epidural  
62278\* Pain management -epidural  
25 62289\* Pain management -epidural  
62298\* Pain management -epidural

*Sites.lst* file

30 ;"ALL" MUST BE FIRST

ALL\* All Centers\*  
aaa\* AA Center\* daniel  
aab\* CenterA\* helen  
35 aac\* CenterB\* debbie  
aad\* CenterC\* jones  
aae\* CenterD\* shannon  
aaf\* Z\* jennifer

40

*Stage1.lst* file

;Last Updated: 04/14/2000

45

DOP\*  
PAYOR\*  
DISPOSITIO\*  
RECTIME\*  
SURGTIME\*

INTTIME\*  
AGE\*  
  
5           ;Anesthesia\*Anesthesia  
  
;-----  
;Miscellaneous (PATINT2)  
;INTTIME2\*IIf(TimeDiff([START],[ENDTIME])<0,0,TimeDiff([START],[ENDTIME]))  
10 Disp3\*IIf(DISPOSITIO="3",True,False)  
Disp2\*IIf(DISPOSITIO="2",True,False)  
  
;-----  
;Miscellaneous (MEDREC)  
15 Anesthesia3\*Left(Anesthesia,3)  
PAYOR3\*Left(PAYOR,3)  
PV\*IIf(pvl="Y",True,False)  
PR\*IIf(prl="Y",True,False)  
Ind2Threshold\*  
  
20 ;-----  
;General Indicators (PATINT2)  
Nummer1\_1\*IIf(Pain\_Complic\_Nausea\_1 OR Pain\_Complic\_Vomiting\_1 OR Pain\_Complic\_InabilityToVoid\_1 OR  
25 Pain\_Complic\_Bleeding\_1 OR Pain\_Complic\_InstabVitalSigns\_1 OR Pain\_Complic\_LevelOfConscChanges\_1 OR  
Pain\_Complic\_RespirProblems\_1,True,False)  
Nummer6\_1\*IIf(Not(IIf(Postop\_Pat\_Int\_Complic\_Nausea\_1 OR Postop\_Pat\_Int\_Complic\_Vomiting\_1 OR  
Postop\_Pat\_Int\_Complic\_Fever\_1 OR Postop\_Pat\_Int\_Complic\_ProblemUrine\_1 OR  
Postop\_Pat\_Int\_Complic\_Bleeding\_1 OR Postop\_Pat\_Int\_Complic\_SignsOfInf\_1,True,False)) AND  
Not(IsNull(PI\_IDN)),True,False)  
30 Nummer7\_1\*IIf((PATINT2.inspain="Y") And Not(folm="Y") And Not(folcom="Y")) And (usem="Y" Or  
inscom="Y") And (folm="Y" Or folcom="Y") And (Left(relief,5)="compl") And (phome="Y"),True,False)  
Denom7\_1\*IIf(phome="Y" And Not(Not(folm="Y") And Not(folcom="Y")),True,False)  
Nummer8\_1\*IIf((qregadm="Excellent" OR qregadm="N/A") And (qpreadm="Excellent" OR qpreadm="N/A") And  
35 (qrecov="Excellent" OR qrecov="N/A") AND (NOT (qregadm="N/A" AND qpreadm="N/A" AND  
qrecov="N/A"))),True,False)  
Nummer9\_1\*IIf(Left(prob,1)="Y" And Left(whocall,1)="Y" And Left(med,1)="Y" And  
Left(app,1)="Y",True,False)  
Nummer10\_1\*IIf(Left(inf,1)="Y" And Left(prob,1)="Y" And Left(whocall,1)="Y" And Left(med,1)="Y" And  
Left(app,1)="Y",True,False)  
  
40 ;-----  
;Pain and Complications (MEDREC)  
Pain\_Complic\_Pain\_1\*IIf(LEFT(PAIN1,1)="Y" OR LEFT(PAIN2,1)="Y" OR LEFT(PAIN3,1)="Y",True,False)  
Pain\_Complic\_Nausea\_1\*IIf(LEFT(NAUS1,1)="Y" OR LEFT(NAUS2,1)="Y" OR LEFT(NAUS3,1)="Y",True,False)  
45 Pain\_Complic\_Vomiting\_1\*IIf(LEFT(VOM1,1)="Y" OR LEFT(VOM2,1)="Y" OR LEFT(VOM3,1)="Y",True,False)  
Pain\_Complic\_InabilityToVoid\_1\*IIf(LEFT(INVOID1,1)="Y" OR LEFT(INVOID2,1)="Y" OR  
LEFT(INVOID3,1)="Y",True,False)  
Pain\_Complic\_Bleeding\_1\*IIf(LEFT(MEDREC.BLEED1,1)="Y" OR LEFT(MEDREC.BLEED2,1)="Y" OR  
LEFT(MEDREC.BLEED3,1)="Y",True,False)  
50 Pain\_Complic\_InstabVitalSigns\_1\*IIf(LEFT(IVS1,1)="Y" OR LEFT(IVS2,1)="Y" OR  
LEFT(IVS3,1)="Y",True,False)  
Pain\_Complic\_LevelOfConscChanges\_1\*IIf(LEFT(LOC1,1)="Y" OR LEFT(LOC2,1)="Y" OR  
LEFT(LOC3,1)="Y",True,False)  
Pain\_Complic\_RespirProblems\_1\*IIf(LEFT(RESPI,1)="Y" OR LEFT(RESPI2,1)="Y" OR  
55 LEFT(RESPI3,1)="Y",True,False)  
  
;-----  
;Pain Control Methods (MEDREC)  
Pain\_Control\_Meth\_PrescrGivenOnDischarge\_1\*IIf(LEFT(PPG,1)="Y",True,False)  
60 Pain\_Control\_Meth\_PainContrMethExplOnDischarge\_1\*IIf(LEFT(PCME,1)="Y",True,False)  
  
Pain\_Control\_Meth\_PainVerb\_1\*IIf(LEFT(PV1,1)="Y",True,False)  
Pain\_Control\_Meth\_MedOrdered\_1\*IIf(LEFT(MOL,1)="Y",True,False) AND Pain\_Control\_Meth\_PainVerb\_1  
Pain\_Control\_Meth\_MedAdmin\_1\*IIf(LEFT(MA1,1)="Y",True,False) AND Pain\_Control\_Meth\_PainVerb\_1  
65 Pain\_Control\_Meth\_MedAdminAndRefused\_1\*IIf(LEFT(MA1,1)="R",True,False) AND Pain\_Control\_Meth\_PainVerb\_1  
Pain\_Control\_Meth\_PainRelieved\_1\*IIf(LEFT(PRI,1)="Y",True,False) AND Pain\_Control\_Meth\_PainVerb\_1  
  
;-----  
;After Leaving the Surgery Center (PATINT2)  
70 After\_Leave\_Surgery\_Problems\_Might\_Have\_1\*IIf(Left(Prob,1)="Y",True,False)  
After\_Leave\_Surgery\_Who\_Call\_1\*IIf(Left(Whocall,1)="Y",True,False)  
After\_Leave\_Surgery\_Meds\_To\_Use\_1\*IIf(Left(Med,1)="Y",True,False)  
After\_Leave\_Surgery\_Had\_Appointment\_1\*IIf(Left(App,1)="Y",True,False)  
After\_Leave\_Surgery\_Had\_All\_Info\_1\*IIf(Left(Inf,1)="Y",True,False)  
  
75 ;-----  
;Postoperative Complications (PATINT2)  
Postop\_Pat\_Int\_Complic\_Nausea\_1\*IIf(PATINT2.nausea3="Y" Or PATINT2.nausea4="Y" Or PATINT2.nausea5="Y"  
Or PATINT2.nausea6="Y" Or PATINT2.nausea7="Y",True,False)  
80 Postop\_Pat\_Int\_Complic\_Vomiting\_1\*IIf(PATINT2.vomit3="Y" Or PATINT2.vomit4="Y" Or PATINT2.vomit5="Y" Or  
PATINT2.vomit6="Y" Or PATINT2.vomit7="Y",True,False)



Patent Application  
Attorney Docket No.: 58367.000003

Postop\_Pat\_Int\_Complic\_Fever\_1\*IIf(PATINT2.fever3="Y" Or PATINT2.fever4="Y" Or PATINT2.fever5="Y" Or  
PATINT2.fever6="Y" Or PATINT2.fever7="Y",True,False)  
Postop\_Pat\_Int\_Complic\_ProblemUrine\_1\*IIf(PATINT2.urine3="Y" Or PATINT2.urine4="Y" Or  
PATINT2.urine5="Y" Or PATINT2.urine6="Y" Or PATINT2.urine7="Y",True,False)  
5 Postop\_Pat\_Int\_Complic\_Bleeding\_1\*IIf(PATINT2.bleed3="Y" Or PATINT2.bleed4="Y" Or PATINT2.bleed5="Y" Or  
PATINT2.bleed6="Y" Or PATINT2.bleed7="Y",True,False)  
Postop\_Pat\_Int\_Complic\_SignsOfInf\_1\*IIf(PATINT2.infec3="Y" Or PATINT2.infec4="Y" Or PATINT2.infec5="Y"  
Or PATINT2.infec6="Y" Or PATINT2.infec7="Y",True,False)  
10 ;-----  
;Postoperative Complications (PATINT2) -- Old Version  
;Postop\_Pat\_Int\_Complic\_Nausea\_1\*IIf(Left(Nausea1,1)="Y",True,False)  
;Postop\_Pat\_Int\_Complic\_Vomiting\_1\*IIf(Left(Vomit1,1)="Y",True,False)  
;Postop\_Pat\_Int\_Complic\_Fever\_1\*IIf(Left(Fever1,1)="Y",True,False)  
15 ;Postop\_Pat\_Int\_Complic\_ProblemUrine\_1\*IIf(Left(Urine1,1)="Y",True,False)  
;Postop\_Pat\_Int\_Complic\_Bleeding\_1\*IIf(Left(Patint2.Bleed1,1)="Y",True,False)  
;Postop\_Pat\_Int\_Complic\_SignsOfInf\_1\*IIf(Left(Infec1,1)="Y",True,False)  
20 ;-----  
;Pain Management at Home (PATINT2)  
Pain\_Manag\_Home\_PostopPainAtHome\_1\*IIf(Left(Phome,1)="Y",True,False)  
Pain\_Manag\_Home\_PostopInstrContrPain\_1\*IIf(Left(Phome,1)="Y" AND Left(Inspain,1)="Y",True,False)  
Pain\_Manag\_Home\_ComplWithInstr\_1\*IIf(Left(Phome,1)="Y" AND (Left(Folm,1)="Y" OR Left(Folcom,1)="Y") AND  
25 IsNull(Foloth),True,False)  
;-----  
;Pain Relief at Home for Patients Who Had Pain (PATINT2)  
Pain\_Relief\_Home\_Completely\_1\*IIf(Left(Phome,1)="Y" AND Left(Relief,3)="Com",True,False)  
Pain\_Relief\_Home\_Greatly\_1\*IIf(Left(Phome,1)="Y" AND Left(Relief,3)="Gre",True,False)  
30 Pain\_Relief\_Home\_Somewhat\_1\*IIf(Left(Phome,1)="Y" AND Left(Relief,3)="Som",True,False)  
Pain\_Relief\_Home\_NotRelieved\_1\*IIf(Left(Phome,1)="Y" AND Left(Relief,3)="Not",True,False)  
;-----  
;Perceived Quality in Registr and Admission Process (PATINT2)  
Perceived\_Quality\_Reg\_And\_Adms\_Excellent\_1\*IIf(Left(Qregadm,3)="Exc",True,False)  
Perceived\_Quality\_Reg\_And\_Adms\_Good\_1\*IIf(Left(Qregadm,3)="Goo",True,False)  
Perceived\_Quality\_Reg\_And\_Adms\_Fair\_1\*IIf(Left(Qregadm,3)="Fai",True,False)  
Perceived\_Quality\_Reg\_And\_Adms\_Poor\_1\*IIf(Left(Qregadm,3)="Poo",True,False)  
40 Perceived\_Quality\_Reg\_And\_Adms\_N-A\_1\*IIf(Left(Qregadm,3)="N/A",True,False)  
;-----  
;Perceived Quality at Preadmission Testing (PATINT2)  
Perceived\_Quality\_Preadmis\_Excellent\_1\*IIf(Left(Qpreadm,3)="Exc",True,False)  
Perceived\_Quality\_Preadmis\_Good\_1\*IIf(Left(Qpreadm,3)="Goo",True,False)  
45 Perceived\_Quality\_Preadmis\_Fair\_1\*IIf(Left(Qpreadm,3)="Fai",True,False)  
Perceived\_Quality\_Preadmis\_Poor\_1\*IIf(Left(Qpreadm,3)="Poo",True,False)  
Perceived\_Quality\_Preadmis\_N-A\_1\*IIf(Left(Qpreadm,3)="N/A",True,False)  
;-----  
;Perceived Quality in Recovery stage in the Center (PATINT2)  
Perceived\_Quality\_Rec\_Stage\_Excellent\_1\*IIf(Left(Qrecov,3)="Exc",True,False)  
Perceived\_Quality\_Rec\_Stage\_Good\_1\*IIf(Left(Qrecov,3)="Goo",True,False)  
Perceived\_Quality\_Rec\_Stage\_Fair\_1\*IIf(Left(Qrecov,3)="Fai",True,False)  
50 Perceived\_Quality\_Rec\_Stage\_Poor\_1\*IIf(Left(Qrecov,3)="Poo",True,False)  
Perceived\_Quality\_Rec\_Stage\_N-A\_1\*IIf(Left(Qrecov,3)="N/A",True,False)  
;-----  
;Age Distribution (MEDREC)  
Age\_Distrib\_0-14\_1\*IIf(AGE>0 AND AGE<15,True,False)  
60 Age\_Distrib\_15-24\_1\*IIf(AGE>=15 AND AGE<25,True,False)  
Age\_Distrib\_25-34\_1\*IIf(AGE>=25 AND AGE<35,True,False)  
Age\_Distrib\_35-44\_1\*IIf(AGE>=35 AND AGE<45,True,False)  
Age\_Distrib\_45-54\_1\*IIf(AGE>=45 AND AGE<55,True,False)  
Age\_Distrib\_55-64\_1\*IIf(AGE>=55 AND AGE<65,True,False)  
65 Age\_Distrib\_65-74\_1\*IIf(AGE>=65 AND AGE<75,True,False)  
Age\_Distrib\_75-84\_1\*IIf(AGE>=75 AND AGE<85,True,False)  
Age\_Distrib\_85+\_1\*IIf(AGE>=85 AND AGE<=120,True,False)  
Age\_Distrib\_Tot\_1\*IIf(AGE>0 AND AGE<=120,True,False)  
70 ;-----  
;Recovery Time Distribution (MEDREC)  
RECTIME-0-30\_1\*IIf(RECTIME>0 and RECTIME<30,True,False)  
RECTIME-30-60\_1\*IIf(RECTIME>=30 and RECTIME<60,True,False)  
RECTIME-60-90\_1\*IIf(RECTIME>=60 and RECTIME<90,True,False)  
75 RECTIME-90-120\_1\*IIf(RECTIME>=90 and RECTIME<120,True,False)  
RECTIME-120-150\_1\*IIf(RECTIME>=120 and RECTIME<150,True,False)  
RECTIME-150-180\_1\*IIf(RECTIME>=150 and RECTIME<180,True,False)  
RECTIME-180-210\_1\*IIf(RECTIME>=180 and RECTIME<210,True,False)  
RECTIME-210-240\_1\*IIf(RECTIME>=210 and RECTIME<240,True,False)  
80 RECTIME-240+\_1\*IIf(RECTIME>=240 ,True,False)  
RECTIME\_TOT\_1\*IIf(RECTIME>0 ,True,False)

```
-----  
;Surgery Time Distribution (MEDREC)  
SURGTIME-0-30_1*IIf(SURGTIME>0 and SURGTIME<30,True,False)  
SURGTIME-30-60_1*IIf(SURGTIME>=30 and SURGTIME<60,True,False)  
SURGTIME-60-90_1*IIf(SURGTIME>=60 and SURGTIME<90,True,False)  
SURGTIME-90-120_1*IIf(SURGTIME>=90 and SURGTIME<120,True,False)  
SURGTIME-120-150_1*IIf(SURGTIME>=120 and SURGTIME<150,True,False)  
SURGTIME-150-180_1*IIf(SURGTIME>=150 and SURGTIME<180,True,False)  
SURGTIME-180-210_1*IIf(SURGTIME>=180 and SURGTIME<210,True,False)  
SURGTIME-210-240_1*IIf(SURGTIME>=210 and SURGTIME<240,True,False)  
SURGTIME-240+_1*IIf(SURGTIME>=240 ,True,False)  
SURGTIME_TOT_1*IIf(SURGTIME>0 ,True,False)  
  
Stage2.1st file  
  
-----  
;Header  
TotMR*      Long*      Count(Site)**  
TotPI*      Long*      Count(PI_IDN)**  
  
-----  
;General Indicators (MEDREC)  
Nummer1*    Long*      Count(IIf(Nummer1_1,True,Null))**  
Nummer2*    Long*      Count(IIf(RECTIME>Ind2Threshold,True,Null))**  
Nummer3*    Long*      Count(IIf(Disp3,True,Null))**  
Nummer4*    Long*      Count(IIf(Disp2,True,Null))**  
Nummer5*    Long*      Count(IIf(PV And Not (PR),True,Null))**  
Denom5*     Long*      Count(IIf(PV,True,Null))**  
  
-----  
;General Indicators (PATINT2)  
Nummer6*    Long*      Count(IIf(Nummer6_1,True,Null))**  
Nummer7*    Long*      Count(IIf(Nummer7_1,True,Null))**  
Denom7*     Long*      Count(IIf(Denom7_1,True,Null))**  
Nummer8*    Long*      Count(IIf(Nummer8_1,True,Null))**  
Nummer9*    Long*      Count(IIf(Nummer9_1,True,Null))**  
Nummer10*   Long*      Count(IIf(Nummer10_1,True,Null))**  
  
-----  
;Complications by Payor (MEDREC)  
TotCare*    Long*      Count(IIf(PAYOR="Medicare",True,Null))**  
Care*       Long*      Count(IIf(PAYOR="Medicare" AND Nummer1_1,True,Null))**  
TotAid*     Long*      Count(IIf(PAYOR="Medicaid",True,Null))**  
Aid*        Long*      Count(IIf(PAYOR="Medicaid" AND Nummer1_1,True,Null))**  
TotCom*     Long*      Count(IIf(Payor3="Com",True,Null))**  
Com*        Long*      Count(IIf(Payor3="Com" AND Nummer1_1,True,Null))**  
TotCap*     Long*      Count(IIf(Payor3="Cap",True,Null))**  
Cap*        Long*      Count(IIf(Payor3="Cap" AND Nummer1_1,True,Null))**  
TotUni*     Long*      Count(IIf(Payor3="Uni",True,Null))**  
Uni*        Long*      Count(IIf(Payor3="Uni" AND Nummer1_1,True,Null))**  
TotWor*     Long*      Count(IIf(Payor3="Wor",True,Null))**  
Wor*        Long*      Count(IIf(Payor3="Wor" AND Nummer1_1,True,Null))**  
TotOth*     Long*      Count(IIf(Payor3="Oth",True,Null))**  
Oth*        Long*      Count(IIf(Payor3="Oth" AND Nummer1_1,True,Null))**  
  
-----  
;Complications by Anesthesia (MEDREC)  
TotEpi*     Long*      Count(IIf(Anesthesia3="Epi",True,Null))**  
Epi*        Long*      Count(IIf(Anesthesia3="Epi" AND Nummer1_1,True,Null))**  
TotGen*     Long*      Count(IIf(Anesthesia3="Gen",True,Null))**  
Gen*        Long*      Count(IIf(Anesthesia3="Gen" AND Nummer1_1,True,Null))**  
TotSpi*     Long*      Count(IIf(Anesthesia3="Spi",True,Null))**  
Spi*        Long*      Count(IIf(Anesthesia3="Spi" AND Nummer1_1,True,Null))**  
TotMAC*     Long*      Count(IIf(Anesthesia3="MAC",True,Null))**  
MAC*        Long*      Count(IIf(Anesthesia3="MAC" AND Nummer1_1,True,Null))**  
TotBlock*   Long*      Count(IIf(Anesthesia3="Blo",True,Null))**  
Block*      Long*      Count(IIf(Anesthesia3="Blo" AND Nummer1_1,True,Null))**  
TotTopical* Long*      Count(IIf(Anesthesia3="Top",True,Null))**  
Topical*    Long*      Count(IIf(Anesthesia3="Top" AND Nummer1_1,True,Null))**  
TotLoc*     Long*      Count(IIf(Anesthesia3="Loc",True,Null))**  
Loc*        Long*      Count(IIf(Anesthesia3="Loc" AND Nummer1_1,True,Null))**  
TotIVC*     Long*      Count(IIf(Anesthesia3="IV-",True,Null))**  
IVC*        Long*      Count(IIf(Anesthesia3="IV-" AND Nummer1_1,True,Null))**  
TotOther*   Long*      Count(IIf(Anesthesia3="Oth",True,Null))**  
Other*      Long*      Count(IIf(Anesthesia3="Oth" AND Nummer1_1,True,Null))**  
TotNone*    Long*      Count(IIf(Anesthesia3="Non",True,Null))**  
None*       Long*      Count(IIf(Anesthesia3="Non" AND Nummer1_1,True,Null))**
```

Patent Application  
Attorney Docket No.: 58367.000003

-----  
5 ;Pain Disposition (MEDREC)  
Patient\_Dispos\_RetainedMore3Hrs\* Long\* Count (Iif(DISPOSITIO="1",True,Null))\*\*  
Patient\_Dispos\_Hospital\* Long\* Count (Iif(DISPOSITIO="2",True,Null))\*\*  
Patient\_Dispos\_Reoperated\* Long\* Count (Iif(DISPOSITIO="3",True,Null))\*\*  
Patient\_Dispos\_Normal\* Long\* Count (Iif(DISPOSITIO="4",True,Null))\*\*  
10 -----  
;Pain and Complications (MEDREC)  
Pain\_Complic\_NoPain-NoComplic\* Long\* Count (Iif(Pain\_Complic\_Pain\_1 OR Pain\_Complic\_Nausea\_1  
OR Pain\_Complic\_Vomiting\_1 OR Pain\_Complic\_InabilityToVoid\_1 OR Pain\_Complic\_Bleeding\_1 OR  
15 Pain\_Complic\_InstabVitalSigns\_1 OR Pain\_Complic\_LevelOfConscChanges\_1 OR  
Pain\_Complic\_RespirProblems\_1,Null,True))\*\*  
Pain\_Complic\_Pain\* Long\* Count (Iif(Pain\_Complic\_Pain\_1,True,Null))\*\*  
Pain\_Complic\_Nausea\* Long\* Count (Iif(Pain\_Complic\_Nausea\_1,True,Null))\*\*  
Pain\_Complic\_Vomiting\* Long\* Count (Iif(Pain\_Complic\_Vomiting\_1,True,Null))\*\*  
20 Pain\_Complic\_InabilityToVoid\* Long\* Count (Iif(Pain\_Complic\_InabilityToVoid\_1,True,Null))\*\*  
Pain\_Complic\_Bleeding\* Long\* Count (Iif(Pain\_Complic\_Bleeding\_1,True,Null))\*\*  
Pain\_Complic\_InstabVitalSigns\* Long\* Count (Iif(Pain\_Complic\_InstabVitalSigns\_1,True,Null))\*\*  
Pain\_Complic\_LevelOfConscChanges\* Long\*  
Count (Iif(Pain\_Complic\_LevelOfConscChanges\_1,True,Null))\*\*  
25 Pain\_Complic\_RespirProblems\* Long\* Count (Iif(Pain\_Complic\_RespirProblems\_1,True,Null))\*\*  
-----  
;Pain Control Methods (MEDREC)  
Pain\_Control\_Meth\_PainVerb\* Long\*  
30 Count (Iif(Pain\_Control\_Meth\_PainVerb\_1,True,Null))\*\*  
Pain\_Control\_Meth\_MedOrdered\* Long\*  
Count (Iif(Pain\_Control\_Meth\_MedOrdered\_1,True,Null))\*\*  
Pain\_Control\_Meth\_MedAdmin\* Long\*  
Count (Iif(Pain\_Control\_Meth\_MedAdmin\_1,True,Null))\*\*  
35 Pain\_Control\_Meth\_MedAdminAndRefused\* Long\*  
Count (Iif(Pain\_Control\_Meth\_MedAdminAndRefused\_1,True,Null))\*\*  
Pain\_Control\_Meth\_PainRelieved\* Long\*  
Count (Iif(Pain\_Control\_Meth\_PainRelieved\_1,True,Null))\*\*  
40 Pain\_Control\_Meth\_PrescrGivenOnDischarge\* Long\*  
Count (Iif(Pain\_Control\_Meth\_PrescrGivenOnDischarge\_1,True,Null))\*\*  
Pain\_Control\_Meth\_PainContrMethExplOnDischarge\* Long\*  
Count (Iif(Pain\_Control\_Meth\_PainContrMethExplOnDischarge\_1,True,Null))\*\*  
-----  
45 ;After Leaving the Surgery Center (PATINT2)  
After\_Leave\_Surgery\_Problems\_Might\_Have\* Long\*  
Count (Iif(After\_Leave\_Surgery\_Problems\_Might\_Have\_1,True,Null))\*\*  
After\_Leave\_Surgery\_Who\_Call\* Long\*  
Count (Iif(After\_Leave\_Surgery\_Who\_Call\_1,True,Null))\*\*  
50 After\_Leave\_Surgery\_Meds\_To\_Use\* Long\*  
Count (Iif(After\_Leave\_Surgery\_Meds\_To\_Use\_1,True,Null))\*\*  
After\_Leave\_Surgery\_Had\_Appointment\* Long\*  
Count (Iif(After\_Leave\_Surgery\_Had\_Appointment\_1,True,Null))\*\*  
After\_Leave\_Surgery\_Had\_All\_Info\* Long\*  
55 Count (Iif(After\_Leave\_Surgery\_Had\_All\_Info\_1,True,Null))\*\*  
-----  
;Postoperative Complications (PATINT2)  
Postop\_Pat\_Int\_Complic\_AnyProblem\* Long\* Count (Iif(Postop\_Pat\_Int\_Complic\_Nausea\_1 OR  
60 Postop\_Pat\_Int\_Complic\_Vomiting\_1 OR Postop\_Pat\_Int\_Complic\_Fever\_1 OR Postop\_Pat\_Int\_Complic\_ProblemUrine\_1  
OR Postop\_Pat\_Int\_Complic\_Bleeding\_1 OR Postop\_Pat\_Int\_Complic\_SignsOfInf\_1,True,Null))\*\*  
Postop\_Pat\_Int\_Complic\_Nausea\* Long\*  
Count (Iif(Postop\_Pat\_Int\_Complic\_Nausea\_1,True,Null))\*\*  
65 Postop\_Pat\_Int\_Complic\_Vomiting\* Long\*  
Count (Iif(Postop\_Pat\_Int\_Complic\_Vomiting\_1,True,Null))\*\*  
Postop\_Pat\_Int\_Complic\_Fever\* Long\*  
Count (Iif(Postop\_Pat\_Int\_Complic\_Fever\_1,True,Null))\*\*  
Postop\_Pat\_Int\_Complic\_ProblemUrine\* Long\*  
70 Count (Iif(Postop\_Pat\_Int\_Complic\_ProblemUrine\_1,True,Null))\*\*  
Postop\_Pat\_Int\_Complic\_Bleeding\* Long\*  
Count (Iif(Postop\_Pat\_Int\_Complic\_Bleeding\_1,True,Null))\*\*  
Postop\_Pat\_Int\_Complic\_SignsOfInf\* Long\*  
Count (Iif(Postop\_Pat\_Int\_Complic\_SignsOfInf\_1,True,Null))\*\*  
75 -----  
;Pain Management at Home (PATINT2)  
Pain\_Manag\_Home\_PostopPainAtHome\* Long\*  
Count (Iif(Pain\_Manag\_Home\_PostopPainAtHome\_1,True,Null))\*\*  
80 Pain\_Manag\_Home\_PostopInstrContrPain\* Long\*  
Count (Iif(Pain\_Manag\_Home\_PostopInstrContrPain\_1,True,Null))\*\*  
-----

Patent Application  
Attorney Docket No.: 58367.000003

```

Pain_Manag_Home_ComplWithInstr*      Long*
Count (IIf (Pain_Manag_Home_ComplWithInstr_1,True,Null))**

5  ;Pain Relief at Home for Patients Who Had Pain (PATINT2)
Pain_Relief_Home_Completely*          Long*      Count (IIf (Pain_Relief_Home_Completely_1,True,Null))**
Pain_Relief_Home_Greatly*             Long*      Count (IIf (Pain_Relief_Home_Greatly_1,True,Null))**
Pain_Relief_Home_Somewhat*            Long*      Count (IIf (Pain_Relief_Home_Somewhat_1,True,Null))**
10 Pain_Relief_Home_NotRelieved*        Long*      Count (IIf (Pain_Relief_Home_NotRelieved_1,True,Null))**

;-----
;Perceived Quality in Registr and Admission Process (PATINT2)
Perceived_Quality_Reg_And_Admis_Excellent* Long*
15 Count (IIf (Perceived_Quality_Reg_And_Admis_Excellent_1,True,Null))**
Perceived_Quality_Reg_And_Admis_Good*      Long*
Count (IIf (Perceived_Quality_Reg_And_Admis_Good_1,True,Null))**
Perceived_Quality_Reg_And_Admis_Fair*      Long*
Count (IIf (Perceived_Quality_Reg_And_Admis_Fair_1,True,Null))**
20 Perceived_Quality_Reg_And_Admis_Poor*     Long*
Count (IIf (Perceived_Quality_Reg_And_Admis_Poor_1,True,Null))**
Perceived_Quality_Reg_And_Admis_N-A*       Long*      Count (IIf ([Perceived_Quality_Reg_And_Admis_N-
A_1],True,Null))**

;-----
25 ;Perceived Quality at Preadmission Testing (PATINT2)
Perceived_Quality_Preadmis_Excellent*     Long*
Count (IIf (Perceived_Quality_Preadmis_Excellent_1,True,Null))**
30 Perceived_Quality_Preadmis_Good*        Long*
Count (IIf (Perceived_Quality_Preadmis_Good_1,True,Null))**
Perceived_Quality_Preadmis_Fair*          Long*
Count (IIf (Perceived_Quality_Preadmis_Fair_1,True,Null))**
Perceived_Quality_Preadmis_Poor*          Long*
Count (IIf (Perceived_Quality_Preadmis_Poor_1,True,Null))**
35 Perceived_Quality_Preadmis_N-A*         Long*      Count (IIf ([Perceived_Quality_Preadmis_N-
A_1],True,Null))**

;-----
40 ;Perceived Quality in Recovery stage in the Center (PATINT2)
Perceived_Quality_Rec_Stage_Excellent*     Long*
Count (IIf (Perceived_Quality_Rec_Stage_Excellent_1,True,Null))**
Perceived_Quality_Rec_Stage_Good*          Long*
Count (IIf (Perceived_Quality_Rec_Stage_Good_1,True,Null))**
45 Perceived_Quality_Rec_Stage_Fair*        Long*
Count (IIf (Perceived_Quality_Rec_Stage_Fair_1,True,Null))**
Perceived_Quality_Rec_Stage_Poor*          Long*
Count (IIf (Perceived_Quality_Rec_Stage_Poor_1,True,Null))**
50 Perceived_Quality_Rec_Stage_N-A*         Long*      Count (IIf ([Perceived_Quality_Rec_Stage_N-
A_1],True,Null))**

;-----
55 ;Age distribution (MEDREC)
Age_Distrib_Avg*      Single*      Avg (IIf (AGE>0 AND AGE<120,AGE,Null))*      Age_Distrib_Tot*
Age_Distrib_0-14*     Long*      Count (IIf ([Age_Distrib_0-14_1],True,Null))**
Age_Distrib_15-24*    Long*      Count (IIf ([Age_Distrib_15-24_1],True,Null))**
Age_Distrib_25-34*    Long*      Count (IIf ([Age_Distrib_25-34_1],True,Null))**
Age_Distrib_35-44*    Long*      Count (IIf ([Age_Distrib_35-44_1],True,Null))**
Age_Distrib_45-54*    Long*      Count (IIf ([Age_Distrib_45-54_1],True,Null))**
Age_Distrib_55-64*    Long*      Count (IIf ([Age_Distrib_55-64_1],True,Null))**
60 Age_Distrib_65-74*    Long*      Count (IIf ([Age_Distrib_65-74_1],True,Null))**
Age_Distrib_75-84*    Long*      Count (IIf ([Age_Distrib_75-84_1],True,Null))**
Age_Distrib_85+*      Long*      Count (IIf ([Age_Distrib_85+_1],True,Null))**
Age_Distrib_Tot*      Long*      Count (IIf ([Age_Distrib_Tot_1],True,Null))**

;-----
65 ;Recovery Time Distribution (MEDREC)
Rectime_Avg*          Single*      Avg (RECTIME)*      RECTIME_TOT*
RECTIME-0-30*         Long*      Count (IIf ([RECTIME-0-30_1],True,Null))**
RECTIME-30-60*        Long*      Count (IIf ([RECTIME-30-60_1],True,Null))**
70 RECTIME-60-90*        Long*      Count (IIf ([RECTIME-60-90_1],True,Null))**
RECTIME-90-120*       Long*      Count (IIf ([RECTIME-90-120_1],True,Null))**
RECTIME-120-150*      Long*      Count (IIf ([RECTIME-120-150_1],True,Null))**
RECTIME-150-180*      Long*      Count (IIf ([RECTIME-150-180_1],True,Null))**
RECTIME-180-210*      Long*      Count (IIf ([RECTIME-180-210_1],True,Null))**
75 RECTIME-210-240*      Long*      Count (IIf ([RECTIME-210-240_1],True,Null))**
RECTIME-240+*         Long*      Count (IIf ([RECTIME-240+_1],True,Null))**
RECTIME_TOT*          Long*      Count (IIf ([RECTIME_TOT_1],True,Null))**

;-----
80 ;Surgery Time Distribution (MEDREC)
Surgtime_Avg*         Single*      Avg (SURGTIME)*      SURGTIME_TOT*
SURGTIME-0-30*        Long*      Count (IIf ([SURGTIME-0-30_1],True,Null))**

```

5      SURGTIME-30-60\*      Long\*      Count(IIf({SURGTIME-30-60\_1},True,Null))\*  
SURGTIME-60-90\*      Long\*      Count(IIf({SURGTIME-60-90\_1},True,Null))\*  
SURGTIME-90-120\*      Long\*      Count(IIf({SURGTIME-90-120\_1},True,Null))\*  
SURGTIME-120-150\*      Long\*      Count(IIf({SURGTIME-120-150\_1},True,Null))\*  
SURGTIME-150-180\*      Long\*      Count(IIf({SURGTIME-150-180\_1},True,Null))\*  
SURGTIME-180-210\*      Long\*      Count(IIf({SURGTIME-180-210\_1},True,Null))\*  
SURGTIME-210-240\*      Long\*      Count(IIf({SURGTIME-210-240\_1},True,Null))\*  
SURGTIME-240\*      Long\*      Count(IIf({SURGTIME-240\_1},True,Null))\*  
10      SURGTIME\_TOT\*      Long\*      Count(IIf({SURGTIME\_TOT\_1},True,Null))\*  
  
;-----  
;Miscellaneous (PATINT2)  
IntTime\_Avg\*      Single\*      Avg(IIf(INTIME>=1 AND INTIME<=20,INTIME,Null))\*      TotPI\*

15      *Corporate\_Members.lst* file

;Format:  
;  
20      ;---Group's Name\*Group's Username\* Group's UserCode\*Allow  
separate members to access their reports (Yes|No)  
;center1  
;center2

25      ---ASC Group\*pinewood\*zaa\*no  
aba  
abh  
acd  
30      acc  
abi  
abj  
abg  
abk

35      *new-soix.ini* file

;This file includes paths to Program folder, INI folder and  
Log folder.  
40      ;Edit it and copy to %windir% directory  
;Caution: Do not put "\" at the end of folder names

;-----Shared parameters  
INIPath  
45      =c:\SOIX\Soix\_Report\_System\INI  
LogPath  
=c:\SOIX\Soix\_Report\_System\Log  
  
OMS2ArchiveDirectory  
50      =c:\SOIX\Soix\_Report\_System\DATA\OMS2\_Archive

OMS2BackupDirectory  
=c:\SOIX\Soix\_Report\_System\DATA\OMS2\_Backup

MDBFile  
5 =c:\SOIX\Soix\_Report\_System\DATA\SOIX.MDB  
LSTPath  
=c:\SOIX\Soix\_Report\_System\lst  
TemplateDirectory  
=c:\SOIX\Soix\_Report\_System\TEMPLATE  
10 UploadDirectory  
=C:\SOIX\WebSites\SOIX\upload  
InternetDirectory  
=C:\SOIX\WebSites\SOIX\Centers  
NewReportsInternetDirectory =C:\SOIX\WebSites\SOIX\NEW  
15 ;-----for paper reports  
SavePathForPaperReport  
=c:\SOIX\WebSites\SOIX\Paper\_Reports  
;Target = Web | Folder  
20 Target =Folder

;-----New sites preparation  
NTSecDirectory =c:\Admin\_Stuff\NTSec  
25 ApacheUsersFile.. =C:\SOIX\WebSites\soix\_users  
ApacheGroupsFile  
=C:\SOIX\WebSites\soix\_groups  
NewCenterTemplateFolder  
=c:\SOIX\Soix\_Report\_System\Template\NewCenterTemplateFolder  
30 PrepareUploadStuff =Yes  
PrepareDownloadStuff =Yes  
PrepareHTMLFiles =Yes

35

EXAMPLE 2

Exemplary indicators for benchmarking ambulatory  
5 surgical procedures in one exemplary embodiment of the  
invention include the following:

INDICATOR 1

General Criteria: Patients experiencing complications of  
surgery during the perioperative period.

10 Indicator Logic And Calculation Concepts

Definition

The percent of all patients experiencing one or more of the  
following complications during the intraoperative or  
postoperative period while in the center:

15 Nausea

Vomiting

Instability of vital signs

Respiratory problems

Level of consciousness changes

20 Hemorrhage/bleeding

Inability to void requiring catheterization (excludes  
intraoperative catheterization)

This indicator may be reported by the payor, the procedure,  
and the anesthesia method

25 Type of Indicator

This is a rate-based indicator of operative complications addressing a process of care with an optimal value of zero. This indicator is stated in terms of a negative outcome; therefore, higher rates should be viewed as opportunities for further review and improvement. Lower rates are generally considered better outcomes.

#### Rationale

Development of operative and postoperative complications may or may not suggest a quality problem. Complications add additional cost to the treatment of patients and/or increase patient risk, length of stay, and recovery time.

#### Calculation Concepts

Numerator = Total Number of Patients Experiencing Complications

Denominator = Total Number of Ambulatory Surgical Patients

Indicator Rate = (Numerator / Denominator) \* 100

#### Data Source

#### Medical Record Abstract Form:

A response under the column for intra or post-operative PROBLEMS for any of the following: nausea, vomiting, inability to void, hemorrhage/bleeding, instability of vital signs, respiratory problems, or LOC changes, will be counted as meeting criteria 1, Patients experiencing complications of surgery during the perioperative period.



This indicator is stated in terms of a negative outcome; therefore, higher rates should be viewed as opportunities for further review and improvement. Lower rates are generally considered better outcomes.

5 Rationale

An additional surgical procedure during the current surgical period may or may not indicate a quality problem. Unplanned returns to surgery subject patients to the additional risks inherent in surgical procedures and the inconvenience or unexpected hardships associated with extended length of stay or prolonged recovery periods. Unplanned returns to surgery also add significantly to the cost of treatment.

Calculation Concepts

Numerator = Total Number of Patients Returned to Surgery

15 Denominator = Total Number of Ambulatory Surgical Patients

Indicator Rate =  $(\text{Numerator} / \text{Denominator}) * 100$

Data Source

Medical Record Abstract Form; patient disposition

INDICATOR 4

20 Patients admitted to the hospital following surgery

Indicator Logic And Calculation Concepts

Definition

Percent of all patients who were admitted to the hospital as inpatients during the intraoperative or postoperative period.

#### Type of Indicator

- 5 This is a rate-based indicator of total admissions following surgery within the postoperative period addressing a process and outcome of care with an optimal value of zero.

This indicator is stated in terms of a negative outcome; therefore, higher rates should be viewed as opportunities

- 10 for further review and improvement. Lower rates are generally considered better outcomes.

#### Rationale

- Admission to the hospital following ambulatory surgery may or may not indicate a quality problem. Although all patients
- 15 included in this study were registered and scheduled as ambulatory surgery patients, admissions following surgery may have been anticipated by the physician in advance (e.g., admissions due to the possibility of additional procedures being performed, the condition of the patient, the time for
- 20 which the procedure was scheduled, etc.). However, unplanned admission subjects a patient to inconvenience or unexpected hardships associated with extended length of stay. Unplanned admissions also add significantly to the cost of treatment.

#### Calculation Concepts

Definition

Percent of all patients interviewed (post-op) who reported  
none of the following problems:

Nausea

5 Vomiting

Fever

Difficulty urinating

Bleeding from the site of the procedure

Excessive redness, swelling, or other sign of infection

10 AND/OR

If one of the above problems was reported, none of the  
following actions were needed to deal with the problem(s)  
they experienced:

Visited the physician

15 Were admitted to the hospital

Received a new or changed prescription

Underwent further surgery

Went to the Emergency Room

Received care from home health care worker

20 Type of indicator

This is a rate-based indicator for patients reporting no  
significant problems during the postoperative period which  
required follow-up care with an optimal value of one.

This indicator is stated in terms of a positive outcome; therefore, lower rates should be viewed as opportunities for further review and improvement. Higher rates are generally considered better outcomes.

5 Rationale

Problems experienced by the patients after discharge related to the surgical procedure may or may not indicate a quality problem. Unexpected problems subject patients to additional risks, inconvenience, or the unexpected hardships associated with prolonged recovery periods. Unexpected problems also add to the cost of treatment.

Calculation Concepts

Numerator = . . . Total Number of Patients Reporting No Problems  
Requiring Action After Discharge.

Denominator = Total Number of Ambulatory Surgical Patients Who  
Participated in the Telephone Interview

Indicator Rate = (Numerator / Denominator) \* 100

Data Source

15 Patient Telephone Interview Form:

In order to be counted as meeting this criterion, a "no" response must be present indicating the patient experienced no nausea, vomiting, fever, etc. after leaving the center AND no boxes checked to indicate medical intervention (i.e.

called doctor, new prescription, etc.). A "yes" response to any of the post-discharge problems WITH medical intervention will not be counted as meeting this criterion.

INDICATOR 7

- 5 Patients expressing pain after discharge who had relief of pain after utilizing pain control methods as instructed.

Indicator Logic And Calculation Concepts

Definition

- Percent of patients interviewed who reported that they were  
10 bothered by pain (related to the surgery) after discharge who:

Reported having instructions for using medicines or for using other comfort measures to control pain

AND

- 15 Reported that they followed the instructions for using medicines and/or other comfort measures

AND

Reported that the methods that they were instructed to use completely relieved their pain.

- 20 Type of Indicator

This is a rate-based indicator of total patients who expressed pain after discharge who were relieved of pain using methods prescribed addressing a process and outcome of care with an optimal value of 100%.

This indicator is stated in terms of a positive outcome; therefore, lower rates should be viewed as opportunities for further review and improvement. Higher rates are generally considered better outcomes.

5 Rationale

Unrelieved post-discharge pain may or may not indicate a quality problem. Such pain, if unrelieved, subjects patients to discomfort or other unexpected hardships associated with pain. Severe or unrelieved pain can result in visits to the physician, ER visits, or prolonged recovery periods, and may add to the dissatisfaction of patients with their treatment.

Calculation Concepts

Numerator = Total Number of Patients  
Completely Relieved of Pain  
Using Prescribed Methods

Denominator = Total Number of Ambulatory  
Surgical Patients Reporting  
Post-Discharge Pain

Indicator Rate = 
$$\frac{(\text{Numerator} / \text{Denominator})}{100} \times$$

15 Data Source

Patient Telephone Interview Form:

Question #3: A response of "yes" must be present in all subcategories a - d., AND

Question #4 must indicate a score of 1 - 5 for "completely relieved." Scores of 6 - 7 = "greatly relieved; " 8 - 9 = "somewhat relieved;" and 10 = "not relieved." Scores of 6 - 10 are not considered as meeting this criterion.

5 INDICATOR 8

Patients satisfied with pre-operative, intraoperative, and postoperative care.

Indicator Logic And Calculation Concepts

Definition

10 Percent of patients interviewed who stated that the quality of care that they received during:

The registration and admission process

AND

..

During the pre-admission testing

15 AND

During the recovery period in the hospital was excellent.

Type of Indicator

This is a rate-based indicator of total patients rating care  
20 as excellent addressing an outcome of care with an optimal value of 100%.

This indicator is stated in terms of a positive outcome; therefore, lower rates should be viewed as opportunities for

and/or "good" are not considered as meeting this criterion.

Responses of "N/A" are ignored and are not counted as either meeting or not meeting the criteria. However, if all three responses are "N/A" the case is considered as NOT meeting

5 the criterion of Indicator 8.

INDICATOR 9.

Patients who received and understood discharge instructions.

Indicator Logic And Calculation Concepts

Definition

10 Percent of patients interviewed who stated that after leaving the hospital they knew:

What problems they might have after surgery

AND ..

Who to call if they had a problem

15 AND

What medicines or other methods to use to control pain

AND

When to see their doctor or when to schedule an appointment.

20 Type of Indicator

This is a rate-based indicator of total patients who received and understood defined aspects of the discharge instructions primarily addressing an outcome of care with an optimal value of 100%.



## Rationale

Patients who do not receive or do not understand their discharge instructions may or may not indicate a quality problem at the hospital level. However, understanding of instructions facilitates a smoother recovery and reduces the inconvenience or unexpected hardships associated with inadequate discharge information. Compliance with discharge plans can reduce patient recovery time.

## Calculation Concepts

Numerator = Total Number of Patients Who Received and Understood Their Discharge Instructions

Denominator = Total Number of Ambulatory Surgical Patients Who Participated in the Telephone Interview

Indicator Rate = 
$$\frac{(\text{Numerator} / \text{Denominator})}{100} \times 100$$

## Data Source

### Patient Telephone Interview Form:

A response of "yes" must be present for each of the first four questions asked under question #1 in order to be considered as meeting this criterion. Responses of "somewhat" are not considered as meeting the criteria.

INDICATOR 10

Patients adequately prepared for self-care at home after discharge.

Indicator Logic And Calculation Concepts

5 Definition

Percent of patients interviewed who stated that after they were home they felt that they had all of the information that they needed to care for themselves

AND

10 that after leaving the hospital they knew

What problems they might have after surgery

AND

Who to call if they had a problem

AND

15 What medicines or other methods to use to control pain

AND

When to see their doctor or when to schedule an appointment.

Type of Indicator

20 This is a rate-based indicator of total patients prepared for care at home after discharge primarily addressing an outcome of care with an optimal value of 100%.

This indicator is stated in terms of a positive outcome; therefore, lower rates should be viewed as opportunities for

further review and improvement. Higher rates are generally considered better outcomes.

#### Rationale

Patients unprepared to care for themselves after discharge  
5 may or may not indicate a quality problem. Understanding of  
self-care instructions facilitates a smoother recovery and  
reduces the inconvenience or unexpected hardships associated  
with inadequate discharge information. Compliance with self-  
care plans can reduce patient recovery time. Lack of  
10 preparation for selfcare adds to the cost of care as  
patients tend to seek care from physicians or hospitals.

#### Calculation Concepts

Numerator =	..	Total Number of Patients Prepared for Self-Care after Discharge
Denominator =		Total Number of Ambulatory Surgical Patients Who Participated in the Telephone Interview
Indicator Rate =		(Numerator / Denominator)*100

#### Data Source

15 Patient Telephone Interview Form:

A response of "yes" must be present to the fifth question  
asked under question #1, AND all of the elements for  
criterion 9 (i.e. "yes" response to the first four

questions) must also be satisfied in order to be considered as meeting criterion 10.

These indicators are exemplary and many can be modified to adjust to changing user needs, new developments in the field and regulatory requirements for example.

For the example given, the first five indicators are displayed as "negative" outcomes (optimal outcome is 0 percent); the last five indicators are displayed as "positive" outcomes with an optimal value of 100 percent.

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the present invention, in addition to those described herein, will be apparent to those of ordinary skill in the art from the foregoing description and accompanying drawings. Thus, such modifications are intended to fall within the scope of the following appended claims. Further, although the present invention has been described herein in the context of a particular implementation in a particular environment for a particular purpose, those of ordinary skill in the art will recognize that its usefulness is not limited thereto and that the present invention can be beneficially implemented in any number of environments for any number of purposes.

Accordingly, the claims set forth below should be construed

in view of the full breath and spirit of the present  
invention as disclosed herein.